Liste modèle des médicaments essentiels

Amitriptyline

Section	Indications
Medicines for other common symptoms in palliative care ◦ Oral > Liquid: 25 mg per 5 mL ◦ Oral > Solid > tablet: 25 mg; 10 mg; 75 mg	Palliative care

Cyclizine

Section	Indications
Medicines for other common symptoms in palliative care ◦ Parenteral > General injections > unspecified: 50 mg per mL (lactate) ◦ Oral > Solid > tablet: 50 mg (hydrochloride)	Palliative care

Dexamethasone

Section	Indications
Medicines for other common symptoms in palliative care ◦ Oral > Liquid: 2 mg per 5 mL (as sodium phosphate); 0.5 mg per 5 mL (as sodium phosphate) ◦ Parenteral > General injections > unspecified: 4 mg per mL in 1 mL ampoule (dexamethasone phosphate (as sodium phosphate) equialent to 3.3 mg dexamethasone base) ◦ Oral > Solid > tablet: 2 mg (as dexamethasone base); 4 mg (as dexamethasone base); 0.5 mg (as dexamethasone base); 0.75 mg (as dexamethasone base); 1.5 mg (as dexamethasone base)	Palliative care

Diazepam

Section	Indications
Medicines for other common symptoms in palliative care ◦ Oral > Liquid: 2 mg per 5 mL ◦ Local > Rectal > Gel: 5 mg per mL in 0.5 mL rectal delivery system; 5 mg per mL in 2 mL rectal delivery system; 5 mg per mL in 4 mL rectal delivery system ◦ Local > Rectal > Solution: 2 mg per mL in 1.25 mL rectal tube; 2 mg per mL in 2.5 mL rectal tube; 4 mg per mL in 2.5 mL rectal tube ◦ Parenteral > General injections > unspecified: 5 mg per mL ◦ Oral > Solid > tablet: 5 mg (scored); 10 mg (scored); 2 mg (scored)	Palliative care

Docusate sodium

Section	Indications
Medicines for other common symptoms in palliative care ◦ Oral > Liquid: 50 mg per 5 mL; 12.5 mg per 5 mL ◦ Oral > Solid > capsule: 100 mg	Palliative care

Dolasetron

Section	Indications
Medicines for other common symptoms in palliative care	Therapeutic equivalent to ondansetron pour Palliative care

Fluoxetine

Section	Indications
Medicines for other common symptoms in palliative care ◦ Oral > Solid: 20 mg (as hydrochloride)	Palliative care

Granisetron

Section	Indications
Medicines for other common symptoms in palliative care	Therapeutic equivalent to ondansetron pour Palliative care

Haloperidol

Section	Indications
Medicines for other common symptoms in palliative care ◦ Oral > Liquid: 2 mg per mL ◦ Oral > Solid: 0.5 mg; 2 mg; 5 mg ◦ Parenteral > General injections > unspecified: 5 mg per mL in 1-mL ampoule	Palliative care

Hyoscine butylbromide

Section	Indications
Medicines for other common symptoms in palliative care ◦ Parenteral > General injections > unspecified: 20 mg per mL injection	Palliative care

Hyoscine hydrobromide

Section	Indications
Medicines for other common symptoms in palliative care ◦ Local > Topical > Transdermal patch: 1 mg per 72 hours transdermal patch ◦ Parenteral > General injections > unspecified: 400 µg per mL injection; 600 µg per mL injection	Palliative care

Lactulose

Section	Indications
Medicines for other common symptoms in palliative care ◦ Oral > Liquid: 3.3 to 3.4 g per 5 mL oral liquid	Palliative care

Loperamide

Section	Indications
Medicines for other common symptoms in palliative care ◦ Oral > Solid > dosage form: 2 mg	Palliative care

Metoclopramide

Section	Indications
Medicines for other common symptoms in palliative care ◦ Oral > Liquid: 5 mg per 5 mL ◦ Oral > Solid: 10 mg (hydrochloride) ◦ Parenteral > General injections > unspecified: 5 mg per mL in 2 mL ampoule (hydrochloride)	Palliative care

Midazolam

Section	Indications
Medicines for other common symptoms in palliative care ◦ Oral > Liquid: 2 mg per mL ◦ Parenteral > General injections > unspecified: 1 mg per mL in 5 mL vial; 5 mg per mL in 1 mL vial; 5 mg per mL in 3 mL vial ◦ Local > Buccal > Solution: ; 5 mg per mL in 0.5 mL pre-filled syringe (for oromucosal administration); 5 mg per mL in 1 mL pre-filled syringe (for oromucosal administration); 5 mg per mL in 1.5 mL pre-filled syringe (for oromucosal administration); 5 mg per mL in 2 mL pre-filled syringe (for oromucosal administration); 10 mg per mL in 0.25 mL pre-filled syringe (for oromucosal administration); 10 mg per mL in 0.5 mL pre-filled syringe (for oromucosal administration); 10 mg per mL in 0.75 mL pre-filled syringe (for oromucosal administration); 10 mg per mL in 1 mL pre-filled syringe (for oromucosal administration) ◦ Oral > Solid > dosage form: 7.5 mg; 15 mg	Palliative care

Ondansetron

Section	Indications
Medicines for other common symptoms in palliative care ◦ Parenteral > General injections > IV: 2 mg per mL in 2 mL ampoule (as hydrochloride dihydrate); 2 mg per mL in 4 mL ampoule (as hydrochloride dihydrate) ◦ Oral > Liquid: 4 mg per 5 mL (as hydrochloride dihydrate) ◦ Oral > Solid: 4 mg (as hydrochloride dihydrate); 8 mg (as hydrochloride dihydrate)	Palliative care

Palonosetron

Section	Indications
Medicines for other common symptoms in palliative care	Therapeutic equivalent to ondansetron pour Palliative care

Senna

Section	Indications
Medicines for other common symptoms in palliative care ◦ Oral > Liquid: 7.5 mg per 5 mL oral liquid	Palliative care

Tropisetron

Section	Indications
Medicines for other common symptoms in palliative care	Therapeutic equivalent to ondansetron pour Palliative care