# Liste modèle des médicaments essentiels

## Acetylcysteine

| Section | Indications |
|---|---|
| Antidotes and other substances used in poisonings > Specific<br>◦ Parenteral > General injections > IV: 200 mg per mL in 10 mL ampoule<br>◦ Oral > Liquid: 10%; 20% | Acute or subacute hepatic failure<br>Exposure to or harmful effects of undetermined intent of analgesics, antipyretics or nonsteroidal anti-inflammatory drugs |

## Activated charcoal

| Section | Indications |
|---|---|
| Antidotes and other substances used in poisonings > Non-specific<br>◦ Oral > Liquid: 50 mg granules for oral suspension | Unspecified injury, poisoning or certain other consequences of external causes |

## Atropine

| Section | Indications |
|---|---|
| Antidotes and other substances used in poisonings > Specific<br>◦ Parenteral > General injections > unspecified: 1 mg in 1 mL ampoule or vial (sulfate) | Exposure to or harmful effects of undetermined intent of pesticides |

## Calcium gluconate

| Section | Indications |
|---|---|
| Antidotes and other substances used in poisonings > Specific<br>◦ Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule or vial (10%) | Hypermagnesaemia |

## Deferoxamine

| Section | Indications |
|---|---|
| Antidotes and other substances used in poisonings > Specific<br>◦ Parenteral > General injections > unspecified: 500 mg in vial powder for injection (mesilate) | Harmful effects of drugs, medicaments or biological substances, not elsewhere classified |

## Dimercaprol

| Section | Indications |
| --- | --- |
| Antidotes and other substances used in poisonings > Specific<br>◦ Parenteral > General injections > IM: 50 mg per mL in 2 mL ampoule; 100 mg per mL in 3 mL ampoule | Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified |

## Fomepizole

| Section | Indications |
| --- | --- |
| Antidotes and other substances used in poisonings > Specific<br>◦ Parenteral > General injections > IV: 5 mg per mL in 20 mL ampoule or vial (sulfate); 1 g per mL in 1.5 mL ampoule or vial (base) | Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified |

## Ipecacuanha

| Section | Indications |
| --- | --- |
| Antidotes and other substances used in poisonings > Non-specific<br>◦ Oral > Liquid: 0.14% ipecacuanha alkaloids calculated as emetine | Unspecified injury, poisoning or certain other consequences of external causes |

## Methionine

| Section | Indications |
| --- | --- |
| Antidotes and other substances used in poisonings > Specific<br>◦ Oral > Solid: 250 mg (DL) | Exposure to or harmful effects of undetermined intent of analgesics, antipyretics or nonsteroidal anti-inflammatory drugs |

## Methylthioninium chloride

| Section | Indications |
| --- | --- |
| Antidotes and other substances used in poisonings > Specific<br>◦ Parenteral > General injections > IV: 10 mg per mL in 10 mL ampoule | Acquired methaemoglobinaemia |

## Naloxone

| Section | Indications |
| --- | --- |
| Antidotes and other substances used in poisonings > Specific<br>◦ Parenteral > General injections > unspecified: 400 µg per 1 mL in ampoule (naloxone hydrochloride) | Unintentional exposure to or harmful effects of opioids or related analgesics |

## Penicillamine

| Section | Indications |
| --- | --- |
| Antidotes and other substances used in poisonings > Specific<br>◦ Oral > Solid: 250 mg | Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified |

## Potassium ferric hexacyanoferrate (Prussian blue)

| Section | Indications |
| --- | --- |
| Antidotes and other substances used in poisonings > Specific<br>◦ Oral > Solid: Powder for oral administration | Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified |

## Pralidoxime

| Section | Indications |
| --- | --- |
| Antidotes and other substances used in poisonings > Specific<br>◦ . | Exposure to or harmful effects of undetermined intent of pesticides |

## Sodium calcium edetate

| Section | Indications |
| --- | --- |
| Antidotes and other substances used in poisonings > Specific<br>◦ Parenteral > General injections > IV: 200 mg per mL in 5 mL ampoule | Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified |

## Sodium nitrite

| Section | Indications |
| --- | --- |
| Antidotes and other substances used in poisonings > Specific<br>◦ Parenteral > General injections > IV: 30 mg per mL in 10 mL ampoule | Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified |

## Sodium thiosulfate

| Section | Indications |
| --- | --- |
| Antidotes and other substances used in poisonings > Specific<br>◦ Parenteral > General injections > IV: 250 mg per mL in 50 mL ampoule | Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified |

## Succimer

| Section | Indications |
| --- | --- |
| Antidotes and other substances used in poisonings > Specific<br>◦ Oral > Solid: 100 mg | Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified |