



## Liste modèle des médicaments essentiels



### Acetylcysteine [Informations générales](#)

#### Section

Antidotes and other substances used in poisonings > Specific

- Parenteral > General injections > IV: 200 mg per mL in 10 mL ampoule
- Oral > Liquid: 10%; 20%

#### Indications

[Acute or subacute hepatic failure](#) [Exposure to or harmful effects of undetermined intent of analgesics, antipyretics or nonsteroidal anti-inflammatory drugs](#)



### Atropine [Informations générales](#)

#### Section

Antidotes and other substances used in poisonings > Specific

- Parenteral > General injections > unspecified: 1 mg in 1 mL ampoule or vial (sulfate)

#### Indications

[Exposure to or harmful effects of undetermined intent of pesticides](#)



### Calcium gluconate [Informations générales](#)

#### Section

Antidotes and other substances used in poisonings > Specific

- Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule or vial (10%)

#### Indications

[Hypermagnesaemia](#)



### Deferoxamine [Informations générales](#)

#### Section

Antidotes and other substances used in poisonings > Specific

- Parenteral > General injections > unspecified: 500 mg in vial powder for injection (mesilate)

#### Indications

[Harmful effects of drugs, medicaments or biological substances, not elsewhere classified](#)



### Dimercaprol [Informations générales](#)

#### Section

Antidotes and other substances used in poisonings > Specific

- Parenteral > General injections > IM: 50 mg per mL in 2 mL ampoule; 100 mg per mL in 3 mL ampoule

#### Indications

[Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified](#)



**Fomepizole** [Informations générales](#)

Section

Antidotes and other substances used in poisonings > Specific

- Parenteral > General injections > IV: 5 mg per mL in 20 mL ampoule or vial (sulfate); 1 g per mL in 1.5 mL ampoule or vial (base)

Indications

[Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified](#)

**Methionine** [Informations générales](#)

Section

Antidotes and other substances used in poisonings > Specific

- Oral > Solid: 250 mg (DL)

Indications

[Exposure to or harmful effects of undetermined intent of analgesics, antipyretics or nonsteroidal anti-inflammatory drugs](#)

**Methylthioninium chloride** [Informations générales](#)

Section

Antidotes and other substances used in poisonings > Specific

- Parenteral > General injections > IV: 10 mg per mL in 10 mL ampoule

Indications

[Acquired methaemoglobinaemia](#)

**Naloxone** [Informations générales](#)

Section

Antidotes and other substances used in poisonings > Specific

- Parenteral > General injections > unspecified: 400 µg per 1 mL in ampoule (naloxone hydrochloride)

Indications

[Unintentional exposure to or harmful effects of opioids or related analgesics](#)

**Penicillamine** [Informations générales](#)

Section

Antidotes and other substances used in poisonings > Specific

- Oral > Solid: 250 mg

Indications

[Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified](#)

**Potassium ferric hexacyanoferrate (Prussian blue)** [Informations générales](#)

Section

Antidotes and other substances used in poisonings > Specific

- Oral > Solid: Powder for oral administration

Indications

[Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified](#)

**Pralidoxime** [Informations générales](#)

Section

Antidotes and other substances used in poisonings > Specific

- .

Indications

[Exposure to or harmful effects of undetermined intent of pesticides](#)

**Sodium calcium edetate** [Informations générales](#)

Section

Antidotes and other substances used in poisonings > Specific

- Parenteral > General injections > IV: 200 mg per mL in 5 mL ampoule

Indications

[Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified](#)

**Sodium nitrite [Informations générales](#)**

Section

Antidotes and other substances used in poisonings > Specific

- Parenteral > General injections > IV: 30 mg per mL in 10 mL ampoule

Indications

[Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified](#)

**Sodium thiosulfate [Informations générales](#)**

Section

Antidotes and other substances used in poisonings > Specific

- Parenteral > General injections > IV: 250 mg per mL in 50 mL ampoule

Indications

[Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified](#)

**Succimer [Informations générales](#)**

Section

Antidotes and other substances used in poisonings > Specific

- Oral > Solid: 100 mg

Indications

[Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified](#)