# Liste modèle des médicaments essentiels

## Cyanocobalamin

| Section | Indications |
| --- | --- |
| Antianaemia medicines<br>◦ . | Anaemias or other erythrocyte disorders |

## Darbepoetin alfa

| Section | Indications |
| --- | --- |
| Antianaemia medicines | Therapeutic equivalent to erythropoiesis-stimulating agents pour<br>Anaemia due to chronic disease |

## Epoetin alfa

| Section | Indications |
| --- | --- |
| Antianaemia medicines | Therapeutic equivalent to erythropoiesis-stimulating agents pour<br>Anaemia due to chronic disease |

## Epoetin beta

| Section | Indications |
| --- | --- |
| Antianaemia medicines | Therapeutic equivalent to erythropoiesis-stimulating agents pour<br>Anaemia due to chronic disease |

## Epoetin theta

| Section | Indications |
| --- | --- |
| Antianaemia medicines | Therapeutic equivalent to erythropoiesis-stimulating agents pour<br>Anaemia due to chronic disease |

## Erythropoiesis-stimulating agents

| Section | Indications |
| --- | --- |
| Antianaemia medicines<br>◦ Parenteral > General injections > unspecified: 1000 IU per 0.5 mL pre-filled syringe; 2000 IU per 0.5 mL pre-filled syringe; 3000 IU per 0.3 mL pre-filled syringe; 4000 IU per 0.4 mL pre-filled syringe; 5000 IU per 0.5 mL pre-filled syringe; 6000 IU per 0.6 mL pre-filled syringe; 8000 IU per 0.8 mL pre-filled syringe; 10000 IU per 1 mL pre-filled syringe; 20000 IU per 0.5 mL pre-filled syringe; 40000 IU per 1 mL pre-filled syringe | Anaemia due to chronic disease |

## Ferrous salt

| Section | Indications |
| --- | --- |
| Antianaemia medicines<br>◦ Oral > Liquid: 25 mg per mL elemental iron; 9 mg per mL elemental iron<br>◦ Oral > Solid > tablet: 60 to 65 mg elemental iron | Iron deficiency anaemia |

## Ferrous salt + folic acid

| Section | Indications |
| --- | --- |
| Antianaemia medicines<br>◦ Oral > Solid: 60 mg iron + 2.8 mg<br>◦ Oral > Solid > tablet: 60 mg elemental iron + 400 µg tablet; 60 mg elemental iron + 2.8 mg | Iron deficiency anaemia<br>Iron deficiency anaemia<br>Spina bifida<br>Clefts of lip, alveolus or palate<br>Anencephaly or similar anomalies |

## Folic acid

| Section | Indications |
| --- | --- |
| Antianaemia medicines<br>◦ Oral > Solid: 400 µg<br>◦ Oral > Liquid: 1 mg per mL<br>◦ Oral > Solid: 1 mg tablet; 5 mg tablet | Cephalocele<br>Anencephaly or similar anomalies<br>Spina bifida<br>Folate deficiency anaemia |

## Hydroxocobalamin

| Section | Indications |
| --- | --- |
| Antianaemia medicines<br>◦ Parenteral > General injections > IM: 1 mg per mL in 1 mL ampoule (as acetate); 1 mg per mL in 1 mL ampoule (as hydrochloride); 1 mg per mL in 1 mL ampoule (as sulfate) | Megaloblastic anaemia due to vitamin B12 deficiency |

## Methoxy polyethylene glycol-epoetin beta

| Section | Indications |
| --- | --- |
| Antianaemia medicines | Therapeutic equivalent to erythropoiesis-stimulating agents pour<br>Anaemia due to chronic disease |