



## Liste modèle des médicaments essentiels



**Cyanocobalamin** [Informations générales](#)

Section  
Antianaemia medicines

Indications

[Anaemias or other erythrocyte disorders](#)



**Darbepoetin alfa** [Informations générales](#)

Section  
Antianaemia medicines

Indications

Therapeutic equivalent to [erythropoiesis-stimulating agents](#) pour [Anaemia due to chronic disease](#)



**Epoetin alfa** [Informations générales](#)

Section  
Antianaemia medicines

Indications

Therapeutic equivalent to [erythropoiesis-stimulating agents](#) pour [Anaemia due to chronic disease](#)



**Epoetin beta** [Informations générales](#)

Section  
Antianaemia medicines

Indications

Therapeutic equivalent to [erythropoiesis-stimulating agents](#) pour [Anaemia due to chronic disease](#)



**Epoetin theta** [Informations générales](#)

Section  
Antianaemia medicines

Indications

Therapeutic equivalent to [erythropoiesis-stimulating agents](#) pour [Anaemia due to chronic disease](#)



**Erythropoiesis-stimulating agents** [Informations générales](#)

Section  
Antianaemia medicines

- Parenteral > General injections > unspecified: 1000 IU per 0.5 mL pre-filled syringe; 2000 IU per 0.5 mL pre-filled syringe; 3000 IU per 0.3 mL pre-filled syringe; 4000 IU per 0.4 mL pre-filled syringe; 5000 IU per 0.5 mL pre-filled syringe; 6000 IU per 0.6 mL pre-filled syringe; 8000 IU per 0.8 mL pre-filled syringe; 10000 IU per 1 mL pre-filled syringe; 20000 IU per 0.5 mL pre-filled syringe; 40000 IU per 1 mL pre-filled syringe

Indications

[Anaemia due to chronic disease](#)



**Ferrous salt** [Informations générales](#)

Section

Antianaemia medicines

- Oral > Liquid: 25 mg per mL elemental iron; 9 mg per mL elemental iron
- Oral > Solid > tablet: 60 to 65 mg elemental iron

Indications

[Iron deficiency anaemia](#)



**Ferrous salt + folic acid** [Informations générales](#)

Section

Antianaemia medicines

- Oral > Solid: 60 mg iron + 2.8 mg
- Oral > Solid > tablet: 60 mg elemental iron + 400 µg tablet; 60 mg elemental iron + 2.8 mg

Indications

[Iron deficiency anaemia](#) [Clefts of lip, alveolus or palate](#) [Iron deficiency anaemia](#) [Anencephaly or similar anomalies](#)

[Spina bifida](#)



**Folic acid** [Informations générales](#)

Section

Antianaemia medicines

- Oral > Solid: 400 µg
- Oral > Liquid: 1 mg per mL
- Oral > Solid: 1 mg tablet; 5 mg tablet

Indications

[Cephalocele](#) [Anencephaly or similar anomalies](#) [Spina bifida](#) [Folate deficiency anaemia](#)



**Hydroxocobalamin** [Informations générales](#)

Section

Antianaemia medicines

- Parenteral > General injections > IM: 1 mg per mL in 1 mL ampoule (as acetate); 1 mg per mL in 1 mL ampoule (as hydrochloride); 1 mg per mL in 1 mL ampoule (as sulfate)

Indications

[Megaloblastic anaemia due to vitamin B12 deficiency](#)



**Methoxy polyethylene glycol-epoetin beta** [Informations générales](#)

Section

Antianaemia medicines

Indications

Therapeutic equivalent to [erythropoiesis-stimulating agents](#) pour [Anaemia due to chronic disease](#)