

Chloroquine

Section	Indications
Antimalarial medicines > Medicines for chemoprevention ◦ Oral > Liquid: 50 mg per 5 mL syrup (as phosphate or sulfate) ◦ Oral > Solid: 150 mg tablet (as phosphate or sulfate)	Malaria due to Plasmodium falciparum Malaria due to Plasmodium ovale Malaria due to Plasmodium malariae Malaria due to Plasmodium vivax
Antimalarial medicines > Medicines for curative treatment ◦ Parenteral > General injections > IV: 40 mg per mL in 5 mL ampoule (as hydrochloride, phosphate or sulfate) ◦ Oral > Liquid: 50 mg per 5 mL syrup (as phosphate or sulfate) ◦ Oral > Solid: 150 mg tablet (as phosphate or sulfate); 100 mg tablet (as phosphate or sulfate) ◦ Oral > Liquid: 50 mg per 5 mL (base) ◦ Oral > Solid > tablet: 150 mg (base)	Malaria due to Plasmodium falciparum Malaria due to Plasmodium vivax
Disease-modifying anti-rheumatic drugs (DMARDs) ◦ Oral > Solid: 100 mg tablet (as phosphate or sulfate); 150 mg tablet (as phosphate or sulfate)	Rheumatoid arthritis
Antimalarial medicines > Medicines for chemoprophylaxis in travellers ◦ Oral > Liquid: 50 mg per 5 mL (base) ◦ Oral > Solid > tablet: 150 mg (base)	Malaria due to Plasmodium vivax

