

Doxycycline

Section

Indications

Access group antibiotics

- Oral > Liquid: 25 mg per 5 mL powder for oral liquid (monohydrate); 50 mg per 5 mL (calcium)
- Parenteral > General injections > unspecified: 100 mg in vial powder for injection
- Oral > Solid > dispersible tablet: 100 mg (as monohydrate)
- Oral > Solid > dosage form: 50 mg (as hyclate); 100 mg (as hyclate)
- Oral > Solid > dosage form: 100 mg (as hyclate)
- Oral > Liquid: 25 mg per 5 mL (anhydrous); 50 mg per 5 mL (anhydrous)
- Oral > Solid: 100 mg (as hyclate); 50 mg (as hyclate)

Antimalarial medicines > Medicines for chemoprophylaxis in travellers

- Oral > Liquid: 50 mg per 5 mL (calcium); 25 mg per 5 mL (monohydrate) powder for oral liquid
- Oral > Solid > dispersible tablet: 100 mg (as monohydrate)
- Oral > Solid > dosage form: 50 mg (as hyclate); 100 mg (as hyclate)

Antimalarial medicines > Medicines for curative treatment

- Oral > Solid: 100 mg (hydrochloride or hyclate); 100 mg dispersible tablet (as monohydrate)

Antimalarial medicines > Medicines for chemoprevention

- Oral > Solid: 100 mg (hydrochloride or hyclate)

First choice

Cholera

Chlamydia trachomatis

Second choice

Bacterial pneumonia (Community-acquired pneumonia - mild to moderate)

Cholera [children]

Chronic obstructive pulmonary disease with acute exacerbation

Other indications

Bacterial infection of unspecified site

Malaria due to Plasmodium falciparum

Malaria due to Plasmodium falciparum

Malaria due to Plasmodium falciparum

