

Acetylsalicylic acid

Section	Indications
Non-opioids and non-steroidal anti-inflammatory medicines (NSAIMs) ◦ Oral > Solid: 100 to 500 mg ◦ Local > Rectal > Suppository: 50 to 150 mg	Pain

Gabapentin

Section	Indications
Non-opioids and non-steroidal anti-inflammatory medicines (NSAIMs) ◦ Oral > Solid: 100 mg; 200 mg; 300 mg; 400 mg; 600 mg; 800 mg	Neuropathic pain

Ibuprofen

Section	Indications
Non-opioids and non-steroidal anti-inflammatory medicines (NSAIMs) ◦ Oral > Liquid: 200 mg per 5 mL; 100 mg per 5 mL ◦ Oral > Solid > tablet: 200 mg; 400 mg; 600 mg	Pain

Paracetamol (acetaminophen)

Section	Indications
Non-opioids and non-steroidal anti-inflammatory medicines (NSAIMs) ◦ Oral > Liquid: 125 mg per 5 mL; 120 mg per 5 mL; 250 mg per 5 mL ◦ Local > Rectal > Suppository: 100 mg; 250 mg ◦ Oral > Solid > dispersible tablet: 100 mg; 250 mg ◦ Oral > Solid > tablet: 250 mg; 325 mg; 500 mg	Pain

