



## Model List of Essential Medicines



### Abacavir [General information](#)

#### Section

Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors

- Oral > Solid: 300 mg tablet (as sulfate)

#### Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)



### Abacavir + dolutegravir + lamivudine [General information](#)

#### Section

Fixed-dose combinations of antiretrovirals

- Oral > Solid > tablet: 60 mg (as sulfate) + 5 mg + 30 mg (dispersible, scored)

#### Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)



### Abacavir + lamivudine [General information](#)

#### Section

Fixed-dose combinations of antiretrovirals

- Oral > Solid: 120 mg (as sulfate) + 60 mg tablet (dispersible, scored)

#### Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)



### Abacavir + lamivudine + lopinavir + ritonavir [General information](#)

#### Section

Fixed-dose combinations of antiretrovirals

- Oral > Solid: 30 mg + 15 mg + 40 mg + 10 mg capsule containing oral granules

#### Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)



### Aciclovir [General information](#)

#### Section

Antiherpes medicines

- Parenteral > General injections > IV: 250 mg in vial powder for solution for infusion (as sodium dihydrate); 25 mg per mL in vial solution for infusion (as sodium)

- Oral > Liquid: 200 mg per 5 mL
- Oral > Solid: 200 mg tablet
- Parenteral > General injections > IV: 250 mg in vial powder for solution for infusion (as sodium dihydrate); 25 mg per mL in vial solution for infusion (as sodium)

Indications

[Herpes simplex infections Zoster Varicella](#)



#### **Atazanavir** [General information](#)

Section

Antiretrovirals > Protease inhibitors

- Oral > Solid: 100 mg (as sulfate); 300 mg (as sulfate)

Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)



#### **Atazanavir + ritonavir** [General information](#)

Section

Antiretrovirals > Protease inhibitors

- Oral > Solid: 300 mg (as sulfate) + 100 mg tablet (heat stable)

Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)



#### **Cobicistat + elvitegravir + emtricitabine + tenofovir disoproxil fumarate** [General information](#)

Section

Fixed-dose combinations of antiretrovirals

- Oral > Solid: 150 mg + 150 mg + 200 mg + 300 mg (tenofovir disoproxil fumarate equivalent to 245 mg tenofovir disoproxil)

Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)



#### **Cobicistat + elvitegravir + emtricitabine+ tenofovir alafenamide** [General information](#)

Section

Fixed-dose combinations of antiretrovirals

- Oral > Solid: 150 mg + 150 mg + 200 mg + 10 mg

Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)



#### **Daclatasvir** [General information](#)

Section

Medicines for hepatitis C > Pangenotypic direct-acting antiviral combinations

- Oral > Solid > tablet: 30 mg (as dihydrochloride); 60 mg (as dihydrochloride)

Indications

[Chronic hepatitis C](#)



#### **Daclatasvir + sofosbuvir** [General information](#)

Section

Medicines for hepatitis C > Pangenotypic direct-acting antiviral combinations

- Oral > Solid > tablet: 60 mg (as dihydrochloride) + 400 mg

Indications

[Chronic hepatitis C](#)



#### **Darunavir** [General information](#)

Section

Antiretrovirals > Protease inhibitors

- Oral > Solid: 75 mg; 400 mg; 600 mg; 800 mg

Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)



**Dasabuvir** [General information](#)

Section

Medicines for hepatitis C > Non-pangenotypic direct-acting antiviral combinations

- Oral > Solid: 250 mg tablet

Indications

[Chronic hepatitis C](#)



**Didanosine** [General information](#)

Section

Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors

- Oral > Liquid: 100 mg buffered powder for oral solution; 167 mg buffered powder for oral solution; 250 mg buffered powder for oral solution
- Oral > Solid: 25 mg buffered chewable dispersible tablet; 50 mg buffered chewable dispersible tablet; 100 mg buffered chewable dispersible tablet; 150 mg buffered chewable dispersible tablet; 200 mg buffered chewable dispersible tablet; 125 mg unbuffered enteric coated capsule; 200 mg unbuffered enteric coated capsule; 250 mg unbuffered enteric coated capsule; 400 mg unbuffered enteric coated capsule

Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)



**Dolutegravir** [General information](#)

Section

Antiretrovirals > Integrase inhibitors

- Oral > Solid: 50 mg tablet; 10 mg tablet (dispersible, scored)

Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)



**Dolutegravir + lamivudine + tenofovir** [General information](#)

Section

Fixed-dose combinations of antiretrovirals

- Oral > Solid: 50 mg + 300 mg + 300 mg (tenofovir disoproxil fumarate equivalent to 245 mg tenofovir disoproxil)

Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)



**Efavirenz** [General information](#)

Section

Antiretrovirals > Non-nucleoside reverse transcriptase inhibitors

- Oral > Solid: 600 mg tablet

Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)



**Efavirenz + emtricitabine + tenofovir** [General information](#)

Section

Fixed-dose combinations of antiretrovirals

- Oral > Solid: 600 mg + 200 mg + 300 mg (tenofovir disoproxil fumarate equivalent to 245 mg tenofovir disoproxil)

Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)



**Efavirenz + lamivudine + tenofovir** [General information](#)

Section

Fixed-dose combinations of antiretrovirals

- Oral > Solid: 400 mg + 300 mg + 300 mg (tenofovir disoproxil fumarate, equivalent to 245 mg tenofovir disoproxil)

Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#) Therapeutic equivalent to [efavirenz + emtricitabine + tenofovir](#) for [Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)



**Elbasvir + grazoprevir** [General information](#)

Section

Medicines for hepatitis C > Non-pangenotypic direct-acting antiviral combinations

- Oral > Solid: 50 mg + 100 mg

Indications

[Chronic hepatitis C](#)



**Emtricitabine** [General information](#)

Section

Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors

- Oral > Liquid: 10 mg per mL
- Oral > Solid: 200 mg

Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)



**Emtricitabine + rilpivirine + tenofovir alafenamide** [General information](#)

Section

Fixed-dose combinations of antiretrovirals

- Oral > Solid: 200 mg + 25 mg + 25 mg

Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)



**Emtricitabine + rilpivirine + tenofovir disoproxil fumarate** [General information](#)

Section

Fixed-dose combinations of antiretrovirals

- Oral > Solid: 200 mg + 25 mg + 300 mg (tenofovir disoproxil fumarate equivalent to 245 mg tenofovir disoproxil)

Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)



**Emtricitabine + tenofovir** [General information](#)

Section

Fixed-dose combinations of antiretrovirals

- Oral > Solid: 200 mg + 300 mg (tenofovir disoproxil fumarate equivalent to 245 mg tenofovir disoproxil)

Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#) [Contact with or exposure to human immunodeficiency virus](#)



**Emtricitabine + tenofovir alafenamide** [General information](#)

Section

Fixed-dose combinations of antiretrovirals

- Oral > Solid: 200 mg + 10 mg; 200 mg + 25 mg

Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)



**Entecavir** [General information](#)

Section

Medicines for hepatitis B > Nucleoside/Nucleotide reverse transcriptase inhibitors

- Oral > Liquid: 0.05 mg per mL
- Oral > Solid: 0.5 mg; 1 mg

Indications

[Chronic hepatitis B](#)



**Glecaprevir + pibrentasvir** [General information](#)

Section

Medicines for hepatitis C > Pangenotypic direct-acting antiviral combinations

- Oral > Solid: 50 mg + 20 mg granules
- Oral > Solid > tablet: 100 mg + 40 mg

Indications

[Chronic hepatitis C](#)



**Indinavir** [General information](#)

Section

Antiretrovirals > Protease inhibitors

- Oral > Solid: 400 mg (as sulfate)

Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)



**Isoniazid + pyridoxine + sulfamethoxazole + trimethoprim** [General information](#)

Section

Medicines for prevention of HIV-related opportunistic infections

- Oral > Solid: 300 mg + 25 mg + 800 mg + 160 mg tablet (scored)

Indications

[Other specified prophylactic measures](#)



**Lamivudine** [General information](#)

Section

Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors

- Oral > Liquid: 50 mg per 5 mL
- Oral > Solid: 150 mg tablet

Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)



**Lamivudine + nevirapine + zidovudine** [General information](#)

Section

Fixed-dose combinations of antiretrovirals

- Oral > Solid: 30 mg + 50 mg + 60 mg Tablet; 150 mg + 200 mg + 300 mg Tablet

Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)



**Lamivudine + tenofovir** [General information](#)

Section

Fixed-dose combinations of antiretrovirals

Indications

Therapeutic equivalent to [emtricitabine + tenofovir](#) for [Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#) Therapeutic equivalent to [emtricitabine + tenofovir](#) for [Contact with or exposure to human immunodeficiency virus](#)



**Lamivudine + zidovudine** [General information](#)

Section

Fixed-dose combinations of antiretrovirals

- Oral > Solid: 30 mg + 60 mg Tablet; 150 mg + 300 mg Tablet

Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)



**Ledipasvir + sofosbuvir** [General information](#)

Section

Medicines for hepatitis C > Non-pangenotypic direct-acting antiviral combinations

- Oral > Solid: 90 mg + 400 mg tablet

Indications

[Chronic hepatitis C](#)



**Lopinavir + ritonavir** [General information](#)

Section

Antiretrovirals > Protease inhibitors

- Oral > Solid > tablet: 100 mg + 25 mg tablet (heat-stable); 200 mg + 50 mg tablet (heat-stable)
- Oral > Solid > dosage form: 40 mg + 10 mg

Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)



**Nelfinavir** [General information](#)

Section

Antiretrovirals > Protease inhibitors

- Oral > Solid: 250 mg (as mesilate)
- Oral > Other: 50 mg per g oral powder

Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)



**Nevirapine** [General information](#)

Section

Antiretrovirals > Non-nucleoside reverse transcriptase inhibitors

- Oral > Liquid: 50 mg per 5 mL
- Oral > Solid > dispersible tablet: 50 mg
- Oral > Solid > tablet: 200 mg

Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)



**Ombitasvir + paritaprevir + ritonavir** [General information](#)

Section

Medicines for hepatitis C > Non-pangenotypic direct-acting antiviral combinations

- Oral > Solid: 12.5 mg + 75 mg + 50 mg tablet

Indications

[Chronic hepatitis C](#)



**Oseltamivir** [General information](#)

Section

Other antivirals

- Oral > Liquid: 6 mg per mL (as phosphate) powder for oral liquid
- Oral > Solid: 30 mg capsule (as phosphate); 45 mg capsule (as phosphate); 75 mg capsule (as phosphate)

Indications

[Influenza due to identified seasonal influenza virus](#)



**Pegylated interferon alfa (2a)** [General information](#)

Section

Medicines for hepatitis C > Other antivirals for hepatitis C

- Parenteral > General injections > SC: 180 µg in vial; 180 µg in prefilled syringe

Indications

[Chronic hepatitis C](#)

**Pegylated interferon alfa (2b) [General information](#)**

## Section

Medicines for hepatitis C > Other antivirals for hepatitis C

- Parenteral > General injections > SC: 80 µg in prefilled syringe; 100 µg in prefilled syringe

## Indications

[Chronic hepatitis C](#)

**Raltegravir [General information](#)**

## Section

Antiretrovirals > Integrase inhibitors

- Oral > Liquid: 100 mg granules for oral suspension
- Oral > Solid: 400 mg tablet; 25 mg tablet (chewable)

## Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)

**Ravidasvir [General information](#)**

## Section

Medicines for hepatitis C > Pangenotypic direct-acting antiviral combinations

- Oral > Solid > tablet: 200 mg

## Indications

[Chronic hepatitis C](#)

**Ribavirin [General information](#)**

## Section

Medicines for hepatitis C > Other antivirals for hepatitis C

- Parenteral > General injections > IV: 1000 mg per 10 mL phosphate buffer solution; 800 mg per 10 mL phosphate buffer solution
- Oral > Solid: 200 mg; 400 mg; 600 mg

## Indications

[Chronic hepatitis C](#)

## Section

Other antivirals

- Parenteral > General injections > IV: 1000 mg per 10 mL phosphate buffer solution; 800 mg per 10 mL phosphate buffer solution
- Oral > Solid: 200 mg; 400 mg; 600 mg

## Indications

[Viral haemorrhagic fever, not elsewhere classified](#)

**Ritonavir [General information](#)**

## Section

Antiretrovirals > Protease inhibitors

- Oral > Solid: 25 mg tablet (heat stable); 100 mg tablet (heat stable)

## Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)

**Saquinavir [General information](#)**

## Section

Antiretrovirals > Protease inhibitors

- Oral > Solid: 200 mg; 500 mg

## Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)

**Simeprevir [General information](#)**

## Section

Medicines for hepatitis C > Non-pangenotypic direct-acting antiviral combinations

- Oral > Solid: 150 mg

Indications

[Chronic hepatitis C](#)



#### **Sofosbuvir** [General information](#)

Section

Medicines for hepatitis C > Pangenotypic direct-acting antiviral combinations

- Oral > Solid > tablet: 200 mg; 400 mg
- Oral > Solid > granules: 200 mg in sachet

Indications

[Chronic hepatitis C](#)



#### **Sofosbuvir + velpatasvir** [General information](#)

Section

Medicines for hepatitis C > Pangenotypic direct-acting antiviral combinations

- Oral > Solid > tablet: 200 mg + 50 mg; 400 mg + 100 mg
- Oral > Solid > granules: 150 mg + 37.5 mg in sachet; 200 mg + 50 mg in sachet

Indications

[Chronic hepatitis C](#)



#### **Stavudine** [General information](#)

Section

Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors

- Oral > Liquid: 5 mg per 5 mL powder for oral solution
- Oral > Solid: 15 mg; 20 mg; 30 mg

Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)



#### **Tenofovir alafenamide** [General information](#)

Section

Medicines for hepatitis B > Nucleoside/Nucleotide reverse transcriptase inhibitors

- Oral > Solid: 25 mg

Indications

[Chronic hepatitis B](#)



#### **Tenofovir disoproxil fumarate** [General information](#)

Section

Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors

- Oral > Solid: 300 mg tablet (equivalent to 245 mg tenofovir disoproxil)

Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#) [Contact with or exposure to human immunodeficiency virus](#)

Section

Medicines for hepatitis B > Nucleoside/Nucleotide reverse transcriptase inhibitors

- Oral > Solid: 300 mg tablet (equivalent to 245 mg tenofovir disoproxil)

Indications

[Chronic hepatitis B](#)



#### **Valganciclovir** [General information](#)

Section

Other antivirals

- Oral > Liquid: 50 mg per mL (as hydrochloride) powder for oral solution
- Oral > Solid > tablet: 450 mg (as hydrochloride)

Indications

[Cytomegaloviral retinitis](#)



#### **Zidovudine** [General information](#)

Section

Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors

- Parenteral > General injections > IV: 10 mg per mL in 20 mL vial solution for IV infusion
- Oral > Liquid: 50 mg per 5 mL
- Oral > Solid > tablet: 300 mg
- Oral > Solid > capsule: 250 mg

Indications

[Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified](#)