

**Amodiaquine****Section**

Antimalarial medicines &gt; Medicines for curative treatment

- Oral > Solid: 153 mg (as hydrochloride); 200 mg (as hydrochloride)

**Indications**Malaria due to *Plasmodium falciparum*Malaria due to *Plasmodium vivax***Amodiaquine - sulfadoxine + pyrimethamine****Section**

Antimalarial medicines &gt; Medicines for chemoprevention

- Oral > Solid > dispersible tablet: 76.5 mg (as hydrochloride) [3] + 250 mg + 12.5 mg [1] in co-package; 153 mg (as hydrochloride) [3] + 500 mg + 25 mg [1] in co-package; 75 mg (as hydrochloride) [3] + 250 mg + 12.5 mg [1] in co-package; 150 mg (as hydrochloride) [3] + 500 mg + 25 mg [1] in co-package

**Indications**

Malaria

**Amphotericin B****Section**

Antileishmaniasis medicines

- Parenteral > General injections > IV: 50 mg in vial powder for injection (liposomal complex); 50 mg in vial powder for injection (as sodium deoxycholate)

**Indications**

Visceral leishmaniasis

**Artemether****Section**

Antimalarial medicines &gt; Medicines for curative treatment

- Parenteral > General injections > IM: 20 mg per mL in 1 mL ampoule (oily injection); 40 mg per mL in 1 mL ampoule (oily injection)

**Indications**Malaria due to *Plasmodium falciparum*Malaria due to *Plasmodium vivax***Artemether + lumefantrine****Section**

Antimalarial medicines &gt; Medicines for curative treatment

- Oral > Solid > dispersible tablet: 20 mg + 120 mg
- Oral > Solid > tablet: 20 mg + 120 mg

**Indications**Malaria due to *Plasmodium falciparum*Malaria due to *Plasmodium vivax*

## Artesunate

### Section

Antimalarial medicines > Medicines for curative treatment

- Parenteral > General injections > IV: 60 mg in vial powder for injection; 30 mg in vial powder for injection; 120 mg in vial powder for injection
- Local > Rectal > Other: 100 mg capsules

### Indications

Malaria due to Plasmodium falciparum

Malaria due to Plasmodium vivax

## Artesunate + amodiaquine

### Section

Antimalarial medicines > Medicines for curative treatment

- Oral > Solid > tablet: 25 mg + 67.5 mg; 50 mg + 135 mg; 100 mg + 270 mg
- Oral > Solid > tablet: 25 mg + 67.5 mg; 50 mg + 135 mg; 100 mg + 270 mg tablet

### Indications

Malaria due to Plasmodium falciparum

Malaria due to Plasmodium vivax

## Artesunate + mefloquine

### Section

Antimalarial medicines > Medicines for curative treatment

- Oral > Solid > tablet: 25 mg + 50 mg (as hydrochloride); 100 mg + 200 mg (as hydrochloride)

### Indications

Malaria due to Plasmodium falciparum

Malaria due to Plasmodium vivax

## Artesunate + pyronaridine

### Section

Antimalarial medicines > Medicines for curative treatment

- Oral > Solid > tablet: 60 mg + 180 mg (tetraphosphate)
- Oral > Solid > granules: 20 mg + 60 mg (tetraphosphate)

### Indications

Malaria due to Plasmodium falciparum

Malaria due to Plasmodium vivax

## Artesunate - sulfadoxine + pyrimethamine

### Section

Antimalarial medicines > Medicines for curative treatment

- Oral > Solid > tablet: 50 mg [3] + 500 mg + 25 mg [1] in copackage (scored)

### Indications

Malaria due to Plasmodium falciparum

Malaria due to Plasmodium vivax

## Benznidazole

### Section

American trypanosomiasis

- Oral > Solid > tablet: 100 mg (scored); 50 mg (scored); 12.5 mg

### Indications

Chagas disease

## Chloroquine

| Section                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        | Indications                                                                                                                                                                                                      |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <a href="#">Antimalarial medicines &gt; Medicines for chemoprevention</a> <ul style="list-style-type: none"><li>◦ Oral &gt; Liquid: 50 mg per 5 mL syrup (as phosphate or sulfate)</li><li>◦ Oral &gt; Solid: 150 mg tablet (as phosphate or sulfate)</li></ul>                                                                                                                                                                                                                                                                                | <a href="#">Malaria due to Plasmodium falciparum</a><br><a href="#">Malaria due to Plasmodium ovale</a><br><a href="#">Malaria due to Plasmodium malariae</a><br><a href="#">Malaria due to Plasmodium vivax</a> |
| <a href="#">Antimalarial medicines &gt; Medicines for curative treatment</a> <ul style="list-style-type: none"><li>◦ Parenteral &gt; General injections &gt; IV: 40 mg per mL in 5 mL ampoule (as hydrochloride, phosphate or sulfate)</li><li>◦ Oral &gt; Liquid: 50 mg per 5 mL syrup (as phosphate or sulfate)</li><li>◦ Oral &gt; Solid: 150 mg tablet (as phosphate or sulfate); 100 mg tablet (as phosphate or sulfate)</li><li>◦ Oral &gt; Liquid: 50 mg per 5 mL (base)</li><li>◦ Oral &gt; Solid &gt; tablet: 150 mg (base)</li></ul> | <a href="#">Malaria due to Plasmodium falciparum</a><br><a href="#">Malaria due to Plasmodium vivax</a>                                                                                                          |
| <a href="#">Antimalarial medicines &gt; Medicines for chemoprophylaxis in travellers</a> <ul style="list-style-type: none"><li>◦ Oral &gt; Liquid: 50 mg per 5 mL (base)</li><li>◦ Oral &gt; Solid &gt; tablet: 150 mg (base)</li></ul>                                                                                                                                                                                                                                                                                                        | <a href="#">Malaria due to Plasmodium vivax</a>                                                                                                                                                                  |

## Dihydroartemisinin + piperaquine

| Section                                                                                                                                                                                                                                                                                                                                                                           | Indications                                                                                             |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------|
| <a href="#">Antimalarial medicines &gt; Medicines for curative treatment</a> <ul style="list-style-type: none"><li>◦ Oral &gt; Solid &gt; dispersible tablet: 20 mg + 160 mg (phosphate); 40 mg + 320 mg (phosphate)</li><li>◦ Oral &gt; Solid &gt; tablet: 40 mg + 320 mg (phosphate); 20 mg + 160 mg (phosphate); 60 mg + 480 (phosphate); 80 mg + 640 mg (phosphate)</li></ul> | <a href="#">Malaria due to Plasmodium falciparum</a><br><a href="#">Malaria due to Plasmodium vivax</a> |

## Diloxanide

| Section                                                                                                                                       | Indications                |
|-----------------------------------------------------------------------------------------------------------------------------------------------|----------------------------|
| <a href="#">Antiamoebic and anti giardiasis medicines</a> <ul style="list-style-type: none"><li>◦ Oral &gt; Solid: 500 mg (furoate)</li></ul> | <a href="#">Amoebiasis</a> |

## Doxycycline

### Section

Antimalarial medicines > Medicines for chemoprophylaxis in travellers

- Oral > Liquid: 50 mg per 5 mL (calcium); 25 mg per 5 mL (monohydrate) powder for oral liquid
- Oral > Solid > dispersible tablet: 100 mg (as monohydrate)
- Oral > Solid > dosage form: 50 mg (as hyclate); 100 mg (as hyclate)

Antimalarial medicines > Medicines for curative treatment

- Oral > Solid: 100 mg (hydrochloride or hyclate); 100 mg dispersible tablet (as monohydrate)

Antimalarial medicines > Medicines for chemoprevention

- Oral > Solid: 100 mg (hydrochloride or hyclate)

### Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium falciparum*

## Eflornithine

### Section

Medicines for the treatment of 2nd stage African trypanosomiasis

- Parenteral > General injections > IV: 200 mg per mL in 50 mL bottle (hydrochloride)

### Indications

Gambiense trypanosomiasis

## Fexinidazole

### Section

Medicines for the treatment of 1st stage African trypanosomiasis

- Oral > Solid > tablet: 600 mg
- Oral > Solid: 600 mg

Medicines for the treatment of 2nd stage African trypanosomiasis

- Oral > Solid > tablet: 600 mg
- Oral > Solid: 600 mg

### Indications

Rhodesiense trypanosomiasis

Gambiense trypanosomiasis

Rhodesiense trypanosomiasis

Gambiense trypanosomiasis

## Hydroxychloroquine

### Section

Antimalarial medicines > Medicines for chemoprevention

### Indications

Therapeutic equivalent to chloroquine for

Malaria due to *Plasmodium falciparum*

## Mefloquine

### Section

Antimalarial medicines > Medicines for curative treatment

- Oral > Solid: 250 mg tablet (hydrochloride)

Antimalarial medicines > Medicines for chemoprophylaxis in travellers

- Oral > Solid > tablet: 250 mg (scored) (as hydrochloride)

Antimalarial medicines > Medicines for chemoprevention

- Oral > Solid: 250 mg tablet (hydrochloride)

### Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium falciparum*

## Meglumine antimoniate

### Section

Antileishmaniasis medicines

- Parenteral > General injections > IM: 1.5 g per 5 mL in 5 mL ampoule

### Indications

Visceral leishmaniasis

Mucocutaneous leishmaniasis

Cutaneous leishmaniasis

## Melarsoprol

### Section

Medicines for the treatment of 2nd stage African trypanosomiasis

- Parenteral > General injections > IV: 3.6% in 5 mL ampoule solution (180 mg of active compound)

### Indications

Gambiense trypanosomiasis

Rhodesiense trypanosomiasis

## Metronidazole

### Section

Antiamoebic and anti giardiasis medicines

- Oral > Liquid: 200 mg per 5 mL (as benzoate)
- Parenteral > General injections > unspecified: 500 mg in 100 mL vial
- Oral > Solid > tablet: 200 mg; 250 mg; 400 mg; 500 mg

### Indications

Amoebiasis

## Miltefosine

### Section

Antileishmaniasis medicines

- Oral > Solid: 10 mg; 50 mg

### Indications

Visceral leishmaniasis

Cutaneous leishmaniasis

Mucocutaneous leishmaniasis

## Nifurtimox

### Section

#### American trypanosomiasis

- Oral > Solid > tablet: 30 mg (scored); 120 mg (scored)

#### Medicines for the treatment of 2nd stage African trypanosomiasis

- Oral > Solid > tablet: 30 mg (scored); 120 mg (scored)

### Indications

Chagas disease

Gambiense trypanosomiasis

## Paromomycin

### Section

#### Antileishmaniasis medicines

- Parenteral > General injections > IM: 750 mg paromomycin base (as sulfate)

#### Antiamoebic and anti giardiasis medicines

- Oral > Liquid: 125 mg per 5 mL as sulfate
- Oral > Solid: 250 mg as sulfate

### Indications

Visceral leishmaniasis

Amoebiasis

## Pentamidine

### Section

#### Antileishmaniasis medicines

- Parenteral > General injections > IM: 200 mg (as isetionate); 300 mg (as isetionate)

#### Medicines for the treatment of 1st stage African trypanosomiasis

- Parenteral > General injections > IM: 300 mg in vial (as isetionate) powder for injection

#### Antipneumocystosis and antitoxoplasmosis medicines

- Oral > Solid > tablet: 200 mg (as isethionate); 300 mg (as isethionate)

### Indications

Visceral leishmaniasis

Gambiense trypanosomiasis

Pneumocystosis

## Primaquine

### Section

#### Antimalarial medicines > Medicines for curative treatment

- Oral > Solid > tablet: 7.5 mg (as phosphate); 15 mg (as phosphate)

### Indications

Malaria due to Plasmodium ovale

Malaria due to Plasmodium falciparum

Malaria due to Plasmodium vivax

## Proguanil

### Section

#### Antimalarial medicines > Medicines for chemoprevention

- Oral > Solid: 100 mg tablet (hydrochloride)

### Indications

Malaria due to Plasmodium falciparum

## Pyrimethamine

### Section

Antipneumocystosis and antitoxoplasmosis medicines

- Oral > Solid: 25 mg

### Indications

Toxoplasmosis

## Quinine

### Section

Antimalarial medicines > Medicines for curative treatment

- Parenteral > General injections > IV: 300 mg per mL in 2 mL ampoule (hydrochloride) solution for infusion; 60 mg per mL in 2 mL ampoule (hydrochloride) solution for infusion

### Indications

Malaria due to Plasmodium falciparum

## Sodium stibogluconate

### Section

Antileishmaniasis medicines

- Parenteral > General injections > unspecified: 100 mg per mL in 30 mL vial

### Indications

Visceral leishmaniasis

Mucocutaneous leishmaniasis

Cutaneous leishmaniasis

## Sulfadiazine

### Section

Antipneumocystosis and antitoxoplasmosis medicines

- Oral > Solid: 500 mg

### Indications

Toxoplasmosis

## Sulfadoxine + pyrimethamine

### Section

Antimalarial medicines > Medicines for chemoprevention

- Oral > Solid: 500 mg + 25 mg tablet
- Oral > Solid > dispersible tablet: 250 mg + 12.5 mg tablet

Antimalarial medicines > Medicines for curative treatment

- Oral > Solid: 500 mg + 25 mg tablet

### Indications

Malaria due to Plasmodium falciparum

Malaria

Malaria due to Plasmodium falciparum

Malaria due to Plasmodium vivax

## Sulfamethoxazole + trimethoprim

### Section

Antipneumocystosis and antitoxoplasmosis medicines

- Parenteral > General injections > IV: 80 mg + 16 mg per mL in 5 mL ampoule; 80 mg + 16 mg per mL in 10 mL ampoule
- Oral > Liquid: 200 mg + 40 mg per 5 mL oral liquid
- Oral > Solid: 100 mg + 20 mg tablet; 400 mg + 80 mg tablet; 800 mg + 160 mg tablet
- Oral > Solid > dispersible tablet: 100 mg + 20 mg

### Indications

Pneumocystosis

## Suramin sodium

| Section                                                                                                                                                                    | Indications                            |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------|
| Medicines for the treatment of 1st stage African trypanosomiasis <ul style="list-style-type: none"><li>◦ Parenteral &gt; General injections &gt; IV: 1 g in vial</li></ul> | <div>Rhodesiense trypanosomiasis</div> |

## Tinidazole

| Section                                  | Indications                                                                       |
|------------------------------------------|-----------------------------------------------------------------------------------|
| Antiamoebic and antigiardiasis medicines | Therapeutic equivalent to <a href="#">metronidazole</a> for <div>Amoebiasis</div> |