## Amodiaquine

| Section | Indications |
| --- | --- |
| Antimalarial medicines > Medicines for curative treatment<br>◦ Oral > Solid: 153 mg (as hydrochloride); 200 mg (as hydrochloride) | Malaria due to Plasmodium falciparum<br>Malaria due to Plasmodium vivax |

## Artemether

| Section | Indications |
| --- | --- |
| Antimalarial medicines > Medicines for curative treatment<br>◦ Parenteral > General injections > IM: 20 mg per mL in 1 mL ampoule (oily injection); 40 mg per mL in 1 mL ampoule (oily injection) | Malaria due to Plasmodium falciparum<br>Malaria due to Plasmodium vivax |

## Artemether + lumefantrine

| Section | Indications |
| --- | --- |
| Antimalarial medicines > Medicines for curative treatment<br>◦ Oral > Solid > dispersible tablet: 20 mg + 120 mg<br>◦ Oral > Solid > tablet: 20 mg + 120 mg | Malaria due to Plasmodium falciparum<br>Malaria due to Plasmodium vivax |

## Artesunate

| Section | Indications |
| --- | --- |
| Antimalarial medicines > Medicines for curative treatment<br>◦ Parenteral > General injections > IV: 60 mg in vial powder for injection; 30 mg in vial powder for injection; 120 mg in vial powder for injection<br>◦ Local > Rectal > Other: 100 mg capsules | Malaria due to Plasmodium falciparum<br>Malaria due to Plasmodium vivax |

## Artesunate + amodiaquine

| Section | Indications |
| --- | --- |
| Antimalarial medicines > Medicines for curative treatment<br>◦ Oral > Solid > tablet: 25 mg + 67.5 mg; 50 mg + 135 mg; 100 mg + 270 mg<br>◦ Oral > Solid > tablet: 25 mg + 67.5 mg; 50 mg + 135 mg; 100 mg + 270 mg tablet | Malaria due to Plasmodium falciparum<br>Malaria due to Plasmodium vivax |

## Artesunate + mefloquine

| Section | Indications |
| --- | --- |
| Antimalarial medicines > Medicines for curative treatment<br>◦ Oral > Solid > tablet: 25 mg + 50 mg (as hydrochloride); 100 mg + 200 mg (as hydrochloride) | Malaria due to Plasmodium falciparum<br>Malaria due to Plasmodium vivax |

## Artesunate + pyronaridine

| Section | Indications |
| --- | --- |
| Antimalarial medicines > Medicines for curative treatment<br>◦ Oral > Solid > tablet: 60 mg + 180 mg (tetraphosphate)<br>◦ Oral > Solid > granules: 20 mg + 60 mg (tetraphosphate) | Malaria due to Plasmodium falciparum<br>Malaria due to Plasmodium vivax |

## Artesunate - sulfadoxine + pyrimethamine

| Section | Indications |
| --- | --- |
| Antimalarial medicines > Medicines for curative treatment<br>◦ Oral > Solid > tablet: 50 mg [3] + 500 mg + 25 mg [1] in copackage (scored) | Malaria due to Plasmodium falciparum<br>Malaria due to Plasmodium vivax |

## Chloroquine

| Section | Indications |
| --- | --- |
| Antimalarial medicines > Medicines for curative treatment<br>◦ Parenteral > General injections > IV: 40 mg per mL in 5 mL ampoule (as hydrochloride, phosphate or sulfate)<br>◦ Oral > Liquid: 50 mg per 5 mL syrup (as phosphate or sulfate)<br>◦ Oral > Solid: 150 mg tablet (as phosphate or sulfate); 100 mg tablet (as phosphate or sulfate)<br>◦ Oral > Liquid: 50 mg per 5 mL (base)<br>◦ Oral > Solid > tablet: 150 mg (base) | Malaria due to Plasmodium falciparum<br>Malaria due to Plasmodium vivax |

## Dihydroartemisinin + piperaquine

| Section | Indications |
| --- | --- |
| Antimalarial medicines > Medicines for curative treatment<br>◦ Oral > Solid > dispersible tablet: 20 mg + 160 mg (phosphate); 40 mg + 320 mg (phosphate)<br>◦ Oral > Solid > tablet: 40 mg + 320 mg (phosphate); 20 mg + 160 mg (phosphate); 60 mg + 480 (phosphate); 80 mg + 640 mg (phosphate) | Malaria due to Plasmodium falciparum<br>Malaria due to Plasmodium vivax |

## Doxycycline

| Section | Indications |
| --- | --- |
| Antimalarial medicines > Medicines for curative treatment<br>◦ Oral > Solid: 100 mg (hydrochloride or hyclate); 100 mg dispersible tablet (as monohydrate) | Malaria due to Plasmodium falciparum |

## Mefloquine

| Section | Indications |
| --- | --- |
| Antimalarial medicines > Medicines for curative treatment<br>◦ Oral > Solid: 250 mg tablet (hydrochloride) | Malaria due to Plasmodium falciparum<br>Malaria due to Plasmodium vivax |

## Primaquine

| Section | Indications |
| --- | --- |
| Antimalarial medicines > Medicines for curative treatment<br>◦ Oral > Solid > tablet: 7.5 mg (as phosphate); 15 mg (as phosphate) | Malaria due to Plasmodium ovale<br>Malaria due to Plasmodium falciparum<br>Malaria due to Plasmodium vivax |

## Quinine

| Section | Indications |
| --- | --- |
| Antimalarial medicines > Medicines for curative treatment<br>◦ Parenteral > General injections > IV: 300 mg per mL in 2 mL ampoule (hydrochloride) solution for infusion; 60 mg per mL in 2 mL ampoule (hydrochloride) solution for infusion | Malaria due to Plasmodium falciparum |

## Sulfadoxine + pyrimethamine

| Section | Indications |
| --- | --- |
| Antimalarial medicines > Medicines for curative treatment<br>◦ Oral > Solid: 500 mg + 25 mg tablet | Malaria due to Plasmodium falciparum<br>Malaria due to Plasmodium vivax |