



Model List of Essential Medicines

Abemaciclib [General information](#)

Section

Targeted therapies

Indications

Therapeutic equivalent to [palbociclib](#) for [Other specified malignant neoplasms of breast](#)

Abiraterone [General information](#)

Section

Hormones and antihormones

- Oral > Solid: 250 mg; 500 mg

Indications

[Malignant neoplasms of prostate](#)

Afatinib [General information](#)

Section

Targeted therapies

Indications

Therapeutic equivalent to [erlotinib](#) for [Other specified malignant neoplasms of bronchus or lung](#)

All-trans retinoic acid [General information](#)

Section

Targeted therapies

- Oral > Solid: 10 mg capsule

Indications

[Acute myeloid leukaemia with recurrent genetic abnormalities](#)

Allopurinol [General information](#)

Section

Supportive medicines

- Parenteral > General injections > IV: 500 mg (as sodium) powder for injection
- Oral > Solid: 100 mg; 300 mg

Indications

[Tumour lysis syndrome](#)

Anastrozole [General information](#)

Section

Hormones and antihormones

- Oral > Solid: 1 mg tablet

Indications

[Other specified malignant neoplasms of breast](#) [Malignant neoplasms of breast](#)



Arsenic trioxide [General information](#)

Section

Cytotoxic medicines

- Parenteral > General injections > IV: 1 mg per mL concentrate for solution for infusion; 2 mg per mL concentrate for solution for infusion

Indications

[Acute myeloid leukaemia with recurrent genetic abnormalities](#)



Asparaginase [General information](#)

Section

Cytotoxic medicines

- Powder for injection: 10 000 IU in vial.
- Parenteral > General injections > unspecified: 10000 IU in vial powder for injection

Indications

[Unspecified malignant neoplasms of ill-defined or unspecified sites](#) [Lymphoid leukaemia, not elsewhere classified](#)



Atezolizumab [General information](#)

Section

Immunomodulators

- Parenteral > General injections > IV: 1.2 g per 20 mL concentrate solution for infusion
- Parenteral > General injections > IV: 840 mg per 14 mL in vial concentrate for solution for infusion; 1200 mg per 20 mL in vial concentrate for solution for infusion

Indications

[Other specified malignant neoplasms of bronchus or lung](#) Therapeutic equivalent to [pembrolizumab](#) for [Other specified malignant neoplasms of bronchus or lung](#) Therapeutic equivalent to [pembrolizumab](#) for [Adenocarcinoma of stomach](#) [Hepatocellular carcinoma of liver](#)



Axicabtagene ciloleucel [General information](#)

Section

Antineoplastics and supportive medicines

- Cell suspension for infusion: 2×10^6 anti-CD19 CAR-positive viable T-cells per kg of body weight.

Indications

[Diffuse large B-cell lymphomas](#)



Axitinib [General information](#)

Section

Targeted therapies

- Oral > Solid > tablet: 1 mg; 3 mg; 5 mg

Indications

[Renal cell carcinoma of kidney, except renal pelvis](#)



Azacitidine [General information](#)

Section

Cytotoxic medicines

- Parenteral > General injections > IV: 100 mg in vial powder for injection

Indications

[Myeloid leukaemia](#)



Bendamustine [General information](#)

Section

Cytotoxic medicines

- Parenteral > General injections > unspecified: 45 mg per 0.5 mL; 180 mg per 2 mL

Indications

[Chronic lymphocytic leukaemia or small lymphocytic lymphoma Follicular lymphoma](#)

Bevacizumab [General information](#)

Section

Immunomodulators

- Parenteral > General injections > IV: 25 mg per mL in 4 mL vial concentrate for solution for infusion; 25 mg per mL in 16 mL vial concentrate for solution for infusion

Indications

[Hepatocellular carcinoma of liver](#)

Bicalutamide [General information](#)

Section

Hormones and antihormones

- Oral > Solid: 50 mg

Indications

[Malignant neoplasms of prostate](#)

Binimetinib [General information](#)

Section

Targeted therapies

Indications

Therapeutic equivalent to [trametinib](#) for [Other specified melanoma of skin](#)

Bleomycin [General information](#)

Section

Cytotoxic medicines

- Parenteral > General injections > unspecified: 15000 IU in vial (as sulfate)

Indications

[Other specified malignant neoplasms of the ovary](#) [Germ cell tumour of testis](#) [Kaposi sarcoma of unspecified primary site](#) [Hodgkin lymphoma](#) [Unspecified malignant neoplasms of ill-defined or unspecified sites](#)

Blinatumomab [General information](#)

Section

Immunomodulators

- Parenteral > General injections > IV: 35 µg in vial; 38.5 µg in vial

Indications

[B lymphoblastic leukaemia or lymphoma, not elsewhere classified](#)

Bortezomib [General information](#)

Section

Targeted therapies

- Parenteral > General injections > unspecified: 3.5 mg in vial powder for injection

Indications

[Plasma cell myeloma](#)

Calcium folinate (leucovorin calcium) [General information](#)

Section

Cytotoxic medicines

- Oral > Solid: 15 mg tablet
- Parenteral > General injections > unspecified: 3 mg per mL in 10 mL ampoule
- Parenteral > General injections > unspecified: 3 mg per mL in 10 mL ampoule; 7.5 mg per mL in 2 mL ampoule; 10 mg per mL in 5 mL ampoule
- Oral > Solid > tablet: 15 mg; 5 mg; 25 mg

Indications

[Unspecified malignant neoplasms of ill-defined or unspecified sites](#) [Malignant neoplasm metastasis in large intestine](#) [Burkitt lymphoma including Burkitt leukaemia](#) [Malignant trophoblastic neoplasms of placenta](#) [Osteosarcoma of bone and articular cartilage of unspecified sites](#) [Malignant neoplasms of rectum](#) [Malignant neoplasms of colon](#)

Capecitabine [General information](#)

Section

Cytotoxic medicines

- Oral > Solid > tablet: 150 mg; 500 mg

Indications

[Malignant neoplasm metastasis in large intestine](#) [Malignant neoplasms of rectum](#) [Malignant neoplasms of colon](#)
[Other specified malignant neoplasms of breast](#)



Carboplatin [General information](#)

Section

Cytotoxic medicines

- Parenteral > General injections > IV: 50 mg per 5 mL; 150 mg per 15 mL; 450 mg per 45 mL; 600 mg per 60 mL

Indications

[Other specified malignant neoplasms of bronchus or lung](#) [Other specified gliomas of brain](#) [Malignant neoplasms of nasopharynx](#) [Malignant neoplasms of breast](#) [Osteosarcoma of bone and articular cartilage of unspecified sites](#) [Other specified carcinomas of ovary](#) [Retinoblastoma](#) [Other specified malignant neoplasms of the ovary](#) [Germ cell tumour of testis](#) [Malignant neoplasms of other or ill-defined sites in the lip, oral cavity or pharynx](#) [Malignant neoplasms of kidney, except renal pelvis](#) [Malignant neoplasms of cervix uteri](#) [Malignant neoplasms of hypopharynx](#) [Malignant neoplasms of nasopharynx](#) [Malignant neoplasms of oropharynx](#)



Cemiplimab [General information](#)

Section

Immunomodulators

- Parenteral > General injections > IV: 50 mg per mL in 7 mL vial concentrate for solution for infusion

Indications

Therapeutic equivalent to [pembrolizumab](#) for [Other specified malignant neoplasms of bronchus or lung](#)
Therapeutic equivalent to [pembrolizumab](#) for [Adenocarcinoma of stomach](#) [Other specified malignant neoplasms of bronchus or lung](#)



Chlorambucil [General information](#)

Section

Cytotoxic medicines

- Oral > Solid: 2 mg tablet

Indications

[Unspecified malignant neoplasms of ill-defined or unspecified sites](#) [Chronic lymphocytic leukaemia or small lymphocytic lymphoma](#)



Cisplatin [General information](#)

Section

Cytotoxic medicines

- Parenteral > General injections > unspecified: 10 mg in vial powder for injection; 50 mg in vial powder for injection
- Parenteral > General injections > IV: 10 mg per 10 mL; 20 mg per 20 mL; 50 mg per 50 mL; 100 mg per 100 mL
- Parenteral > General injections > IV: 50 mg per 50 mL; 100 mg per 100 mL; 10 mg per 10 mL; 20 mg per 20 mL

Indications

[Unspecified malignant neoplasms of ill-defined or unspecified sites](#) [Other specified gliomas of brain](#) [Other specified malignant neoplasms of the ovary](#) [Other specified malignant neoplasms of bronchus or lung](#) [Malignant neoplasms of nasopharynx](#) [Germ cell tumour of testis](#) [Malignant neoplasms of other or ill-defined sites in the lip, oral cavity or pharynx](#) [Osteosarcoma of bone and articular cartilage of unspecified sites](#) [Malignant neoplasms of cervix uteri](#) [Malignant neoplasms of hypopharynx](#) [Malignant neoplasms of nasopharynx](#) [Malignant neoplasms of oropharynx](#)



Cladribine [General information](#)

Section

Cytotoxic medicines

- Parenteral > General injections > IV: 1 mg per mL in 10 mL vial; 2 mg per mL in 5 mL vial

Indications

[Langerhans cell histiocytosis](#)



Cobimetinib [General information](#)

Section

Targeted therapies

Indications

Therapeutic equivalent to [trametinib](#) for [Other specified melanoma of skin](#)



Crizotinib [General information](#)

Section

Targeted therapies

- Oral > Solid: 200 mg; 250 mg
- Oral > Solid > capsule: 200 mg; 250 mg

Indications

[Other specified malignant neoplasms of bronchus or lung Anaplastic large cell lymphoma, ALK-positive](#)



Cyclophosphamide [General information](#)

Section

Cytotoxic medicines

- Oral > Solid: 25 mg tablet
- Parenteral > General injections > unspecified: 500 mg in vial powder for injection
- Parenteral > General injections > IV: 500 mg in vial powder for injection; 1 g in vial powder for injection; 2 g in vial powder for injection
- Oral > Solid > dosage form: 25 mg; 50 mg

Indications

[Unspecified malignant neoplasms of ill-defined or unspecified sites Other specified malignant neoplasms of breast Follicular lymphoma Malignant trophoblastic neoplasms of placenta Malignant neoplasms of breast Diffuse large B-cell lymphoma, not otherwise specified Chronic lymphocytic leukaemia or small lymphocytic lymphoma Hodgkin lymphoma Plasma cell myeloma Other specified gliomas of brain Burkitt lymphoma including Burkitt leukaemia Ewing sarcoma of bone and articular cartilage of unspecified sites Malignant neoplasms of kidney, except renal pelvis Lymphoid leukaemia, not elsewhere classified Anaplastic large cell lymphoma, ALK-negative Anaplastic large cell lymphoma, ALK-positive Rhabdomyosarcoma primary site](#)



Cytarabine [General information](#)

Section

Cytotoxic medicines

- Parenteral > General injections > unspecified: 100 mg in vial powder for injection
- Parenteral > General injections > unspecified: 100 mg in vial powder for injection; 100 mg per mL in vial

Indications

[Unspecified malignant neoplasms of ill-defined or unspecified sites Acute myeloid leukaemia with recurrent genetic abnormalities Burkitt lymphoma including Burkitt leukaemia Lymphoid leukaemia, not elsewhere classified Myeloid leukaemia Anaplastic large cell lymphoma, ALK-negative Anaplastic large cell lymphoma, ALK-positive Langerhans cell histiocytosis](#)



Dabrafenib [General information](#)

Section

Targeted therapies

- Oral > Solid: 50 mg; 75 mg

Indications

[Other specified melanoma of skin](#)



Dacarbazine [General information](#)

Section

Cytotoxic medicines

- Parenteral > General injections > IV: 100 mg in vial powder for injection; 200 mg in vial powder for injection

Indications

[Hodgkin lymphoma](#)



Dactinomycin [General information](#)

Section

Cytotoxic medicines

- Parenteral > General injections > IV: 500 µg in vial powder for injection

Indications

[Malignant trophoblastic neoplasms of placenta](#) [Unspecified malignant neoplasms of ill-defined or unspecified sites](#)
[Malignant neoplasms of kidney, except renal pelvis](#) [Rhabdomyosarcoma primary site](#) [Ewing sarcoma of bone and articular cartilage of unspecified sites](#)

Daratumumab [General information](#)

Section

Immunomodulators

- Parenteral > General injections > IV: 100 mg per 5 mL; 400 mg per 20 mL

Indications

[Plasma cell myeloma](#)

Darbepoetin alfa [General information](#)

Section

Supportive medicines

Indications

Therapeutic equivalent to [erythropoiesis-stimulating agents](#) for [Drug-induced aplastic anaemia](#)

Dasatinib [General information](#)

Section

Targeted therapies

- Oral > Solid: 20 mg tablet; 50 mg tablet; 70 mg tablet; 80 mg tablet; 100 mg tablet; 140 mg tablet

Indications

[Chronic myeloid leukaemia, not elsewhere classified](#)

Daunorubicin [General information](#)

Section

Cytotoxic medicines

- Parenteral > General injections > IV: 50 mg in vial (as hydrochloride) powder for injection; 20 mg in vial (as hydrochloride) powder for injection; 2 mg per mL in vial (as hydrochloride); 5 mg per mL in vial (as hydrochloride)

Indications

[Acute myeloid leukaemia with recurrent genetic abnormalities](#) [Lymphoid leukaemia, not elsewhere classified](#)
[Myeloid leukaemia](#)

Dexamethasone [General information](#)

Section

Hormones and antihormones

- Oral > Liquid: 2 mg per 5 mL
- Oral > Solid: 2 mg; 4 mg
- Parenteral > General injections > unspecified: 4 mg per mL in 1 mL ampoule (as disodium phosphate salt)
- Parenteral > General injections > unspecified: 4 mg per mL in 1 mL ampoule dexamethasone phosphate (as sodium phosphate, equivalent to 3.3 mg/mL dexamethasone base)
- Oral > Solid > tablet: 4 mg (as dexamethasone base)
- Oral > Liquid: 2 mg per 5 mL (as sodium phosphate)
- Parenteral > General injections > unspecified: 4 mg per mL in 1 mL ampoule dexamethasone phosphate (as sodium phosphate, equivalent to 3.3 mg/mL dexamethasone base)
- Oral > Solid > tablet: 2 mg (as dexamethasone base); 4 mg (as dexamethasone base)

Indications

[Anaplastic large cell lymphoma, ALK-negative](#) [Plasma cell myeloma](#) [Burkitt lymphoma including Burkitt leukaemia](#)
[Lymphoid leukaemia, not elsewhere classified](#) [Anaplastic large cell lymphoma, ALK-positive](#)

Docetaxel [General information](#)

Section

Cytotoxic medicines

- Parenteral > General injections > IV: 20 mg per mL; 40 mg per mL

Indications

[Other specified malignant neoplasms of breast](#) [Malignant neoplasms of breast](#) [Malignant neoplasms of prostate](#)
[Unspecified malignant neoplasms of ill-defined or unspecified sites](#)

Dostarlimab [General information](#)

Section

Immunomodulators

- Parenteral > General injections > IV: 50 mg per mL in 10 mL vial concentrate for solution for infusion

Indications

[Other specified malignant neoplasms of corpus uteri](#)



Doxorubicin [General information](#)

Section

Cytotoxic medicines

- Parenteral > General injections > IV: 10 mg in vial (hydrochloride); 50 mg in vial (hydrochloride)
- Parenteral > General injections > IV: 10 mg in vial (hydrochloride) powder for injection; 50 mg in vial (hydrochloride) powder for injection; 2 mg per mL in 5 mL vial (hydrochloride); 2 mg per mL in 25 mL vial (hydrochloride)

Indications

[Unspecified malignant neoplasms of ill-defined or unspecified sites](#) [Rhabdomyosarcoma primary site](#) [Follicular lymphoma](#) [Burkitt lymphoma including Burkitt leukaemia](#) [Malignant neoplasms of kidney, except renal pelvis](#) [Malignant neoplasms of breast](#) [Osteosarcoma of bone and articular cartilage of unspecified sites](#) [Lymphoid leukaemia, not elsewhere classified](#) [Diffuse large B-cell lymphoma, not otherwise specified](#) [Other specified malignant neoplasms of breast](#) [Hodgkin lymphoma](#) [Plasma cell myeloma](#) [Ewing sarcoma of bone and articular cartilage of unspecified sites](#) [Anaplastic large cell lymphoma, ALK-negative](#) [Anaplastic large cell lymphoma, ALK-positive](#) [Kaposi sarcoma of unspecified primary site](#)



Doxorubicin (as pegylated liposomal) [General information](#)

Section

Cytotoxic medicines

- Parenteral > General injections > IV: 2 mg per mL in 10 mL vial (hydrochloride); 2 mg per mL in 25 mL vial (hydrochloride)

Indications

[Kaposi sarcoma of unspecified primary site](#)



Durvalumab [General information](#)

Section

Immunomodulators

- Parenteral > General injections > IV: 50 mg per mL in 2.4 mL vial concentrate solution for infusion; 50 mg per mL in 10 mL vial concentrate solution for infusion

Indications

[Other specified malignant neoplasms of bronchus or lung](#) [Hepatocellular carcinoma of liver](#) [Malignant neoplasms of biliary tract, distal bile duct](#) [Malignant neoplasm of intrahepatic bile ducts](#) [Malignant neoplasms of gallbladder](#) [Malignant neoplasms of proximal biliary tract, cystic duct](#)



Encorafenib [General information](#)

Section

Targeted therapies

Indications

Therapeutic equivalent to [dabrafenib](#) for [Other specified melanoma of skin](#)



Enzalutamide [General information](#)

Section

Hormones and antihormones

- Oral > Solid: 40 mg

Indications

Therapeutic equivalent to [abiraterone](#) for [Malignant neoplasms of prostate](#) [Malignant neoplasms of prostate](#)



Epoetin alfa [General information](#)

Section

Supportive medicines

Indications

Therapeutic equivalent to [erythropoiesis-stimulating agents](#) for [Drug-induced aplastic anaemia](#)



Epoetin beta [General information](#)

Section

Supportive medicines

Indications

Therapeutic equivalent to [erythropoiesis-stimulating agents](#) for [Drug-induced aplastic anaemia](#)

Erlotinib [General information](#)

Section

Targeted therapies

- Oral > Solid: 100 mg; 150 mg

Indications

[Other specified malignant neoplasms of bronchus or lung](#)

Erythropoiesis-stimulating agents [General information](#)

Section

Supportive medicines

- Parenteral > General injections > unspecified: 2000 IU per mL in vial (single-dose); 3000 IU per mL in vial (single-dose); 4000 IU per mL in vial (single-dose); 10000 IU per mL in vial (single-dose); 40000 IU per mL in vial (single-dose); 20000 IU per mL in vial (multiple-dose); 20000 IU per 2 mL in vial (multiple-dose)

Indications

[Drug-induced aplastic anaemia](#)

Etoposide [General information](#)

Section

Cytotoxic medicines

- Parenteral > General injections > IV: 20 mg per mL in 5 mL ampoule
- Oral > Solid: 100 mg capsule
- Parenteral > General injections > IV: 20 mg per mL in 5 mL ampoule; 100 mg in vial powder for injection (as phosphate)
- Oral > Solid: 100 mg capsule; 50 mg capsule

Indications

[Unspecified malignant neoplasms of ill-defined or unspecified sites](#) [Burkitt lymphoma including Burkitt leukaemia](#) [Ewing sarcoma of bone and articular cartilage of unspecified sites](#) [Lymphoid leukaemia, not elsewhere classified](#) [Myeloid leukaemia](#) [Other specified malignant neoplasms of the ovary](#) [Other specified malignant neoplasms of bronchus or lung](#) [Germ cell tumour of testis](#) [Malignant trophoblastic neoplasms of placenta](#) [Malignant neoplasms of kidney, except renal pelvis](#) [Osteosarcoma of bone and articular cartilage of unspecified sites](#) [Hodgkin lymphoma](#) [Retinoblastoma](#) [Anaplastic large cell lymphoma, ALK-negative](#) [Anaplastic large cell lymphoma, ALK-positive](#)

Everolimus [General information](#)

Section

Targeted therapies

- Oral > Solid > dispersible tablet: 2 mg; 3 mg; 5 mg
- Oral > Solid > tablet: 2.5 mg; 5 mg; 7.5 mg; 10 mg

Indications

[Other specified gliomas of brain](#)

Filgrastim [General information](#)

Section

Immunomodulators

- Parenteral > General injections > SC: 120 µg per 0.2 mL in prefilled syringe; 300 µg per 0.5 mL in prefilled syringe; 480 µg per 0.8 mL in prefilled syringe; 300 µg per mL in 1 mL vial; 480 µg per 1.6 mL in 1.6 mL vial

Indications

[Acquired neutropaenia](#)

Fludarabine [General information](#)

Section

Cytotoxic medicines

- Parenteral > General injections > IV: 50 mg in vial (fludarabine phosphate) powder for injection
- Oral > Solid: 10 mg tablet (fludarabine phosphate)

Indications

[Chronic lymphocytic leukaemia or small lymphocytic lymphoma](#)



Fluorouracil [General information](#)

Section

Cytotoxic medicines

- Parenteral > General injections > IV: 50 mg per mL in 5 mL ampoule
- Parenteral > General injections > IV: 50 mg per mL in vial

Indications

[Unspecified malignant neoplasms of ill-defined or unspecified sites](#) [Malignant neoplasms of nasopharynx](#) [Malignant neoplasm metastasis in large intestine](#) [Malignant neoplasms of breast](#) [Malignant neoplasms of rectum](#) [Malignant neoplasms of colon](#)



Flutamide [General information](#)

Section

Hormones and antihormones

Indications

Therapeutic equivalent to [bicalutamide](#) for [Malignant neoplasms of prostate](#)



Fulvestrant [General information](#)

Section

Hormones and antihormones

- Parenteral > General injections > IV: 250 mg per 5 mL

Indications

[Malignant neoplasms of breast](#)



Gefitinib [General information](#)

Section

Targeted therapies

Indications

Therapeutic equivalent to [erlotinib](#) for [Other specified malignant neoplasms of bronchus or lung](#)



Gemcitabine [General information](#)

Section

Cytotoxic medicines

- Parenteral > General injections > IV: 200 mg in vial powder for injection; 1 g in vial powder for injection

Indications

[Other specified carcinomas of ovary](#) [Other specified malignant neoplasms of bronchus or lung](#)



Goserelin [General information](#)

Section

Hormones and antihormones

Indications

Therapeutic equivalent to [leuprorelin](#) for [Malignant neoplasms of breast](#) Therapeutic equivalent to [leuprorelin](#) for [Malignant neoplasms of prostate](#)



Hydrocortisone [General information](#)

Section

Hormones and antihormones

- Parenteral > General injections > unspecified: 100 mg in vial (as sodium succinate)

Indications

[Lymphoid leukaemia, not elsewhere classified](#) [Unspecified malignant neoplasms of ill-defined or unspecified sites](#) [Burkitt lymphoma including Burkitt leukaemia](#)



Hydroxycarbamide (hydroxyurea) [General information](#)

Section

Cytotoxic medicines

- Oral > Solid: 200 mg; 250 mg; 300 mg; 400 mg; 500 mg; 1 g
- Oral > Solid > dosage form: 200 mg; 300 mg; 400 mg; 500 mg; 1 g; 100 mg

Indications

[Squamous cell carcinoma of oropharynx](#) [Chronic myeloid leukaemia, not elsewhere classified](#)



Ibrutinib [General information](#)

Section

Targeted therapies

- Oral > Solid: 140 mg

Indications

[Chronic lymphocytic leukaemia or small lymphocytic lymphoma](#)



Ifosfamide [General information](#)

Section

Cytotoxic medicines

- Parenteral > General injections > IV: 500 mg in vial powder for injection; 1 g in vial powder for injection; 2 g in vial powder for injection
- Parenteral > General injections > IV: 1 g in vial powder for injection; 2 g in vial powder for injection

Indications

[Other specified malignant neoplasms of the ovary](#) [Germ cell tumour of testis](#) [Ewing sarcoma of bone and articular cartilage of unspecified sites](#) [Osteosarcoma of bone and articular cartilage of unspecified sites](#) [Rhabdomyosarcoma primary site](#) [Unspecified malignant neoplasms of ill-defined or unspecified sites](#) [Burkitt lymphoma including Burkitt leukaemia](#) [Malignant neoplasms of kidney, except renal pelvis](#) [Anaplastic large cell lymphoma, ALK-negative](#) [Anaplastic large cell lymphoma, ALK-positive](#)



Imatinib [General information](#)

Section

Targeted therapies

- Oral > Solid: 100 mg; 400 mg

Indications

[B lymphoblastic leukaemia or lymphoma with t\(9;22\) \(q34;q11.2\); BCR-ABL1](#) [Chronic myeloid leukaemia, not elsewhere classified](#) [Gastrointestinal stromal tumour of unspecified gastrointestinal sites](#)



Ipilimumab [General information](#)

Section

Immunomodulators

- Parenteral > General injections > IV: 5 mg per mL in 10 mL vial concentrate for solution for infusion; 20 mg per mL in 40 mL vial concentrate for solution for infusion

Indications

[Other specified malignant neoplasms of bronchus or lung](#) [Renal cell carcinoma of kidney, except renal pelvis](#) [Squamous cell carcinoma of oesophagus](#) [Melanoma of skin](#)



Irinotecan [General information](#)

Section

Cytotoxic medicines

- Parenteral > General injections > IV: 40 mg per 2 mL in 2 mL vial; 100 mg per 5 mL in 5 mL vial; 500 mg per 25 mL in 25 mL vial

Indications

[Malignant neoplasm metastasis in large intestine](#) [Malignant neoplasms of kidney, except renal pelvis](#) [Rhabdomyosarcoma primary site](#)



Lenalidomide [General information](#)

Section

Immunomodulators

- Oral > Solid: 25 mg

Indications

[Plasma cell myeloma](#)



Lenvatinib [General information](#)

Section

Targeted therapies

- Oral > Solid > capsule: 4 mg; 10 mg

Indications

[Renal cell carcinoma of kidney, except renal pelvis](#)



Leuprorelin [General information](#)

Section

Hormones and antihormones

- Parenteral > General injections > IM: 7.5 mg in prefilled syringe; 22.5 mg in prefilled syringe

Indications

[Malignant neoplasms of breast](#) [Malignant neoplasms of prostate](#)



Lisocabtagene maraleucel [General information](#)

Section

Antineoplastics and supportive medicines

Indications

Therapeutic equivalent to [axicabtagene ciloleucel](#) for [Diffuse large B-cell lymphomas](#)



Melphalan [General information](#)

Section

Cytotoxic medicines

- Parenteral > General injections > IV: 50 mg in vial powder for injection
- Oral > Solid: 2 mg

Indications

[Plasma cell myeloma](#)



Mercaptopurine [General information](#)

Section

Cytotoxic medicines

- Oral > Solid: 50 mg tablet
- Oral > Liquid: 20 mg per mL

Indications

[Unspecified malignant neoplasms of ill-defined or unspecified sites](#) [Lymphoid leukaemia, not elsewhere classified](#) [Acute myeloid leukaemia with recurrent genetic abnormalities](#) [Langerhans cell histiocytosis](#)



Mesna [General information](#)

Section

Supportive medicines

- Parenteral > General injections > IV: 100 mg per mL in 4 mL ampoule; 100 mg per mL in 10 mL ampoule
- Oral > Solid: 400 mg; 600 mg
- Parenteral > General injections > IV: 100 mg per mL in 4 mL ampoule; 100 mg per mL in 10 mL ampoule; 100 mg per mL in 2 mL ampoule
- Oral > Solid > tablet: 400 mg; 600 mg
- Oral > Other: 600 mg
- Oral > Solid > tablet: 400 mg

Indications

[Unspecified malignant neoplasms of ill-defined or unspecified sites](#) [Osteosarcoma of bone and articular cartilage of other specified sites](#) [Burkitt lymphoma including Burkitt leukaemia](#) [Ewing sarcoma of bone and articular cartilage of unspecified sites](#) [Malignant neoplasms of kidney, except renal pelvis](#) [Other specified malignant neoplasms of the ovary](#) [Germ cell tumour of testis](#) [Rhabdomyosarcoma primary site](#)



Methotrexate [General information](#)

Section

Cytotoxic medicines

- Parenteral > General injections > IV: 50 mg in vial (as sodium salt)
- Oral > Solid: 2.5 mg (as sodium salt)
- Parenteral > General injections > IV: 50 mg in vial (as sodium salt); 50 mg per 2 mL; 1000 mg per 10 mL concentrated injection

Indications

[Unspecified malignant neoplasms of ill-defined or unspecified sites](#) [Lymphoid leukaemia, not elsewhere classified](#)

[Acute myeloid leukaemia with recurrent genetic abnormalities](#) [Burkitt lymphoma including Burkitt leukaemia](#) [Malignant trophoblastic neoplasms of placenta](#) [Malignant neoplasms of breast](#) [Osteosarcoma of bone and articular cartilage of unspecified sites](#) [Anaplastic large cell lymphoma, ALK-negative](#) [Anaplastic large cell lymphoma, ALK-positive](#) [Langerhans cell histiocytosis](#)

Methylprednisolone [General information](#)

Section

Hormones and antihormones

- Parenteral > General injections > IV: 40 mg in vial (as sodium succinate); 125 mg in vial (as sodium succinate)
- Parenteral > General injections > IV: 40 mg in vial (as sodium succinate) powder for injection; 125 mg in vial (as sodium succinate) powder for injection

Indications

[Burkitt lymphoma including Burkitt leukaemia](#) [Lymphoid leukaemia, not elsewhere classified](#)

Nilotinib [General information](#)

Section

Targeted therapies

- Oral > Solid > capsule: 150 mg; 200 mg

Indications

[Chronic myeloid leukaemia, not elsewhere classified](#)

Nilutamide [General information](#)

Section

Hormones and antihormones

Indications

Therapeutic equivalent to [bicalutamide](#) for [Malignant neoplasms of prostate](#)

Nivolumab [General information](#)

Section

Immunomodulators

- Parenteral > General injections > IV: 10 mg per mL concentrate solution for infusion
- Parenteral > General injections > IV: 10 mg per mL in 4 mL vial concentrate solution for infusion; 10 mg per mL in 10 mL vial concentrate solution for infusion; 10 mg per mL in 12 mL vial concentrate solution for infusion; 10 mg per mL in 24 mL vial concentrate solution for infusion
- Parenteral > General injections > IV: 10 mg per mL in 24 mL vial concentrate solution for infusion; 10 mg per mL in 4 mL vial concentrate solution for infusion; 10 mg per mL in 10 mL vial concentrate solution for infusion; 10 mg per mL in 12 mL vial concentrate solution for infusion

Indications

Therapeutic equivalent to [pembrolizumab](#) for [Melanoma of skin](#) [Other specified malignant neoplasms of bronchus or lung](#) [Melanoma of skin](#) [Renal cell carcinoma of kidney, except renal pelvis](#) [Squamous cell carcinoma of oesophagus](#) [Adenocarcinoma of oesophagogastric junction](#) [Adenocarcinoma of stomach](#)

Osimertinib [General information](#)

Section

Targeted therapies

- Oral > Solid: 40 mg (as mesylate); 80 mg (as mesylate)

Indications

[Other specified malignant neoplasms of bronchus or lung](#)

Oxaliplatin [General information](#)

Section

Cytotoxic medicines

- Parenteral > General injections > IV: 50 mg in vial powder for injection; 100 mg in vial powder for injection; 50 mg per 10 mL in 10 mL vial; 100 mg per 20 mL in 20 mL vial; 200 mg per 40 mL in 40 mL vial

Indications

[Malignant neoplasm metastasis in large intestine](#) [Malignant neoplasms of colon](#)

Paclitaxel [General information](#)

Section

Cytotoxic medicines

- Parenteral > General injections > IV: 6 mg per mL

Indications

[Other specified malignant neoplasms of the ovary](#) [Other specified malignant neoplasms of bronchus or lung](#) [Malignant neoplasms of nasopharynx](#) [Kaposi sarcoma of unspecified primary site](#) [Other specified malignant neoplasms of breast](#) [Other specified carcinomas of ovary](#) [Malignant neoplasms of breast](#) [Malignant neoplasms of cervix uteri](#)

Palbociclib [General information](#)

Section

Targeted therapies

- Oral > Solid: 75 mg; 100 mg; 125 mg

Indications

[Other specified malignant neoplasms of breast](#)

Panitumumab [General information](#)

Section

Targeted therapies

- Parenteral > General injections > IV: 20 mg per mL in 5 mL vial concentrate for solution for infusion; 20 mg per mL in 20 mL vial

Indications

[Other specified malignant neoplasms of large intestine](#)

Pegaspargase [General information](#)

Section

Cytotoxic medicines

- Parenteral > General injections > IV: 3750 units per 5 mL in vial; 3750 units in vial powder for injection

Indications

[Lymphoid leukaemia, not elsewhere classified](#)

Pegfilgrastim [General information](#)

Section

Immunomodulators

- Parenteral > General injections > SC: 6 mg per 0.6 mL in pre-filled syringe

Indications

[Acquired neutropaenia](#)

Pembrolizumab [General information](#)

Section

Immunomodulators

- Parenteral > General injections > IV: 25 mg per mL in 4 mL vial
- Parenteral > General injections > IV: 25 mg per mL in 4 mL vial powder for injection
- Parenteral > General injections > IV: 25 mg per mL in 4 mL vial concentrate for solution for infusion

Indications

[Melanoma of skin](#) [Malignant neoplasms of cervix uteri](#) [Other specified malignant neoplasms of large intestine](#) [Other specified malignant neoplasms of bronchus or lung](#) [Adenocarcinoma of stomach](#) [Squamous cell carcinoma of hypopharynx and variants](#) [Squamous cell carcinoma of other or ill-defined sites in the lip, oral cavity or pharynx](#) [Squamous cell carcinoma of oropharynx](#) [Squamous cell carcinoma of nasopharynx](#) [Malignant neoplasms of oesophagogastric junction](#) [Renal cell carcinoma of kidney, except renal pelvis](#) [Carcinoma of breast, specialised type](#) [Squamous cell carcinoma of oesophagus](#) Therapeutic equivalent to [nivolumab](#) for [Melanoma of skin](#)

Pertuzumab [General information](#)

Section

Targeted therapies

- Parenteral > General injections > IV: 240 mg per 14 mL in vial concentrated solution

Indications

[Carcinoma of breast, specialised type](#)

Prednisolone [General information](#)

Section

Hormones and antihormones

- Oral > Liquid: 5 mg per mL
- Oral > Solid: 5 mg; 25 mg

Indications

[Unspecified malignant neoplasms of ill-defined or unspecified sites](#) [Follicular lymphoma](#) [Lymphoid leukaemia, not elsewhere classified](#) [Diffuse large B-cell lymphoma, not otherwise specified](#) [Chronic lymphocytic leukaemia or small lymphocytic lymphoma](#) [Hodgkin lymphoma](#) [Plasma cell myeloma](#) [Malignant neoplasms of prostate](#) [Burkitt lymphoma including Burkitt leukaemia](#) [Anaplastic large cell lymphoma, ALK-negative](#) [Anaplastic large cell lymphoma, ALK-positive](#) [Langerhans cell histiocytosis](#)



Prednisone [General information](#)

Section

Hormones and antihormones

Indications

Therapeutic equivalent to [prednisolone](#) for [Follicular lymphoma](#) Therapeutic equivalent to [prednisolone](#) for [Lymphoid leukaemia, not elsewhere classified](#) Therapeutic equivalent to [prednisolone](#) for [Diffuse large B-cell lymphoma, not otherwise specified](#) Therapeutic equivalent to [prednisolone](#) for [Chronic lymphocytic leukaemia or small lymphocytic lymphoma](#) Therapeutic equivalent to [prednisolone](#) for [Hodgkin lymphoma](#) Therapeutic equivalent to [prednisolone](#) for [Plasma cell myeloma](#) Therapeutic equivalent to [prednisolone](#) for [Malignant neoplasms of prostate](#) Therapeutic equivalent to [prednisolone](#) for [Anaplastic large cell lymphoma, ALK-negative](#) Therapeutic equivalent to [prednisolone](#) for [Anaplastic large cell lymphoma, ALK-positive](#) Therapeutic equivalent to [prednisolone](#) for [Langerhans cell histiocytosis](#)



Procarbazine [General information](#)

Section

Cytotoxic medicines

- Oral > Solid: 50 mg (as hydrochloride)
- Oral > Solid > capsule: 50 mg (as hydrochloride)

Indications

[Unspecified malignant neoplasms of ill-defined or unspecified sites](#) [Hodgkin lymphoma](#)



Rasburicase [General information](#)

Section

Supportive medicines

- Parenteral > General injections > IV: 1.5 mg in vial powder and 1 mL solvent for solution; 7.5 mg in vial powder and 5 mL solvent for solution

Indications

[Tumour lysis syndrome](#)



Realgar-indigo naturalis formulation [General information](#)

Section

Cytotoxic medicines

- Oral > Solid: 270 mg (containing tetra-arsenic tetra-sulfide 30 mg)

Indications

[Acute myeloid leukaemia with recurrent genetic abnormalities](#)



Ribociclib [General information](#)

Section

Targeted therapies

Indications

Therapeutic equivalent to [palbociclib](#) for [Other specified malignant neoplasms of breast](#)



Rituximab [General information](#)

Section

Targeted therapies

- Parenteral > General injections > IV: 100 mg per 10 mL in 10 mL vial; 500 mg per 50 mL in 50 mL vial

Indications

[Follicular lymphoma](#) [Diffuse large B-cell lymphoma, not otherwise specified](#) [Chronic lymphocytic leukaemia or small lymphocytic lymphoma](#) [Burkitt lymphoma including Burkitt leukaemia](#)



Tamoxifen [General information](#)

Section

Hormones and antihormones

- Oral > Solid: 10 mg (as citrate); 20 mg (as citrate)

Indications

[Other specified malignant neoplasms of breast](#) [Malignant neoplasms of breast](#)

Temozolomide [General information](#)

Section

Cytotoxic medicines

- Parenteral > General injections > IV: 100 mg in vial powder for injection
- Oral > Solid > capsule: 5 mg; 20 mg; 100 mg; 140 mg; 180 mg; 250 mg

Indications

[Ewing sarcoma of bone and articular cartilage of unspecified sites](#) [Other specified gliomas of brain](#) [Neuroblastoma](#)
[NOS](#) [Unspecified malignant neoplasms of ill-defined or unspecified sites](#)

Thalidomide [General information](#)

Section

Immunomodulators

- Oral > Solid: 50 mg

Indications

[Plasma cell myeloma](#)

Tioguanine [General information](#)

Section

Cytotoxic medicines

- Oral > Solid: 40 mg

Indications

[Lymphoid leukaemia, not elsewhere classified](#)

Tisagenlecleucel [General information](#)

Section

Antineoplastics and supportive medicines

Indications

Therapeutic equivalent to [axicabtagene ciloleucel](#) for [Diffuse large B-cell lymphomas](#)

Tislelizumab [General information](#)

Section

Immunomodulators

- Parenteral > General injections > IV: 100 mg per 10 mL
- Parenteral > General injections > IV: 100 mg per 10 mL concentrate for solution for infusion
- Parenteral > General injections > IV: 100 mg per 10 mL concentrate for solution for infusion

Indications

[Malignant neoplasms of urinary tract](#) [Hodgkin lymphoma](#) [Other specified malignant neoplasms of bronchus or lung](#)
[Squamous cell carcinoma of oesophagus](#)

Toripalimab [General information](#)

Section

Immunomodulators

- Parenteral > General injections > IV: 240 mg per 6 mL in vial

Indications

[Malignant neoplasms of nasopharynx](#) [Squamous cell carcinoma of oesophagus](#)

Trametinib [General information](#)

Section

Targeted therapies

- Oral > Solid: 0.5 mg; 2 mg

Indications

[Other specified melanoma of skin](#)

Trastuzumab [General information](#)

Section

Targeted therapies

- Parenteral > General injections > IV: 60 mg in vial powder for injection; 150 mg in vial powder for injection; 440 mg in vial powder for injection

Indications

[Carcinoma of breast, specialised type](#)

Trastuzumab emtansine [General information](#)

Section

Targeted therapies

- Parenteral > General injections > IV: 100 mg in vial powder for injection; 160 mg in vial powder for injection

Indications

[Carcinoma of breast, specialised type](#)

Tremelimumab [General information](#)

Section

Immunomodulators

- Parenteral > General injections > IV: 20 mg per mL in 1.25 mL vial concentrate solution for infusion; 20 mg per mL in 15 mL vial concentrate solution for infusion

Indications

[Hepatocellular carcinoma of liver](#) [Other specified malignant neoplasms of bronchus or lung](#)

Triptorelin [General information](#)

Section

Hormones and antihormones

Indications

Therapeutic equivalent to [leuprorelin](#) for [Malignant neoplasms of breast](#) Therapeutic equivalent to [leuprorelin](#) for [Malignant neoplasms of prostate](#)

Vemurafenib [General information](#)

Section

Targeted therapies

Indications

Therapeutic equivalent to [dabrafenib](#) for [Other specified melanoma of skin](#)

Vinblastine [General information](#)

Section

Cytotoxic medicines

- Parenteral > General injections > IV: 10 mg in vial (vinblastine sulfate)
- Parenteral > General injections > IV: 10 mg in vial (vinblastine sulfate); 10 mg per 10 mL in vial (vinblastine sulfate)

Indications

[Unspecified malignant neoplasms of ill-defined or unspecified sites](#) [Other specified gliomas of brain](#) [Hodgkin lymphoma](#) [Other specified malignant neoplasms of the ovary](#) [Germ cell tumour of testis](#) [Kaposi sarcoma of unspecified primary site](#) [Anaplastic large cell lymphoma, ALK-negative](#) [Anaplastic large cell lymphoma, ALK-positive](#) [Langerhans cell histiocytosis](#)

Vincristine [General information](#)

Section

Cytotoxic medicines

- Parenteral > General injections > IV: 1 mg in vial (vincristine sulfate); 5 mg in vial (vincristine sulfate)
- Parenteral > General injections > IV: 1 mg in vial (vincristine sulfate); 5 mg in vial (vincristine sulfate); 1 mg per mL in vial (vincristine sulfate); 2 mg per 2 mL in vial (vincristine sulfate)

Indications

[Unspecified malignant neoplasms of ill-defined or unspecified sites](#) [Other specified gliomas of brain](#) [Follicular lymphoma](#) [Malignant trophoblastic neoplasms of placenta](#) [Diffuse large B-cell lymphoma, not otherwise specified](#)

[Kaposi sarcoma of unspecified primary site](#) [Rhabdomyosarcoma primary site](#) [Hodgkin lymphoma](#) [Burkitt lymphoma including Burkitt leukaemia](#) [Ewing sarcoma of bone and articular cartilage of unspecified sites](#) [Malignant neoplasms of kidney, except renal pelvis](#) [Lymphoid leukaemia, not elsewhere classified](#) [Retinoblastoma](#) [Langerhans cell histiocytosis](#)

Vinorelbine [General information](#)

Section

Cytotoxic medicines

- Parenteral > General injections > IV: 10 mg per mL in 1 mL vial; 10 mg per mL in 5 mL vial
- Parenteral > General injections > IV: 10 mg per mL in 1 mL vial; 10 mg per mL in 5 mL vial
- Oral > Solid > capsule: 20 mg; 30 mg; 80 mg

Indications

[Other specified malignant neoplasms of bronchus or lung](#) [Other specified malignant neoplasms of breast](#) [Rhabdomyosarcoma primary site](#)

Zanubrutinib [General information](#)

Section

Targeted therapies

- Oral > Solid: 80 mg
- Oral > Solid > capsule: 80 mg

Indications

[Mantle cell lymphoma](#) [Chronic lymphocytic leukaemia or small lymphocytic lymphoma](#) Therapeutic equivalent to [ibrutinib](#) for [Chronic lymphocytic leukaemia or small lymphocytic lymphoma](#)

Zoledronic acid [General information](#)

Section

Supportive medicines

- Parenteral > General injections > IV: 4 mg per 5 mL in 5 mL vial; 4 mg per 100 mL in 100 mL bottle

Indications

[Malignant neoplasm metastasis in bone or bone marrow](#)