

Acetylsalicylic acid

Section	Indications
Anti-platelet medicines ◦ Oral > Solid: 100 mg	Acute myocardial infarction Cerebral ischaemic stroke

Acetylsalicylic acid + atorvastatin + ramipril

Section	Indications
Fixed-dose combinations for prevention of atherosclerotic cardiovascular disease ◦ Oral > Solid > tablet: 100 mg + 20 mg + 2.5 mg; 100 mg + 20 mg + 5 mg; 100 mg + 20 mg + 10 mg; 100 mg + 40 mg + 2.5 mg; 100 mg + 40 mg + 5 mg; 100 mg + 40 mg + 10 mg	Atherosclerotic chronic arterial occlusive disease Chronic ischaemic heart disease Cerebral ischaemic stroke

Acetylsalicylic acid + simvastatin + ramipril + atenolol + hydrochlorothiazide

Section	Indications
Fixed-dose combinations for prevention of atherosclerotic cardiovascular disease ◦ Oral > Solid > tablet: 100 mg + 20 mg + 5 mg + 50 mg + 12.5 mg	Atherosclerotic chronic arterial occlusive disease Chronic ischaemic heart disease Cerebral ischaemic stroke

Alteplase

Section	Indications
Thrombolytic medicines ◦ Parenteral > General injections > IV: 10 mg in vial powder for injection; 20 mg in vial powder for injection; 50 mg in vial powder for injection	Cerebral ischaemic stroke

Amiodarone

Section	Indications
Antiarrhythmic medicines ◦ Parenteral > General injections > IV: 50 mg per mL in 3 ampoule (hydrochloride) ◦ Oral > Solid: 100 mg (hydrochloride); 200 mg (hydrochloride); 400 mg (hydrochloride)	Cardiac arrhythmia

Amlodipine

Section

Antihypertensive medicines

- Oral > Solid: 5 mg (as maleate, mesylate or besylate)

Indications

Essential hypertension

Atenolol

Section

Antianginal medicines

- Oral > Solid: 50 mg; 100 mg

Antihypertensive medicines

- Oral > Solid: 50 mg; 100 mg

Indications

Angina pectoris

Essential hypertension

Therapeutic equivalent to [bisoprolol](#) for

Essential hypertension

Atorvastatin

Section

Lipid-lowering agents

Indications

Therapeutic equivalent to [simvastatin](#) for

Mixed hyperlipidaemia

Therapeutic equivalent to [simvastatin](#) for

Coronary atherosclerosis

Atorvastatin + perindopril + amlodipine

Section

Fixed-dose combinations for prevention of atherosclerotic cardiovascular disease

- Oral > Solid > tablet: 20 mg + 5 mg + 5 mg; 20 mg + 10 mg + 10 mg; 40 mg + 5 mg + 5 mg; 40 mg + 10 mg + 10 mg

Indications

Atherosclerotic chronic arterial occlusive disease

Chronic ischaemic heart disease

Cerebral ischaemic stroke

Bisoprolol

Section

Antianginal medicines

- Oral > Solid: 1.25 mg; 5 mg

Antihypertensive medicines

- Oral > Solid: 1.25 mg; 5 mg

Antiarrhythmic medicines

- Oral > Solid: 1.25 mg; 5 mg

Medicines used in heart failure

- Oral > Solid: 1.25 mg; 5 mg

Indications

Angina pectoris

Essential hypertension

Cardiac arrhythmia

Heart failure

Bumetanide

Section

Medicines used in heart failure

Indications

Therapeutic equivalent to [furosemide](#) for

Heart failure

Captopril

Section

Medicines used in heart failure

- Oral > Solid: 25 mg

Indications

Heart failure

Carvedilol

Section

Antianginal medicines

Therapeutic equivalent to [bisoprolol](#) for

Angina pectoris

Antihypertensive medicines

Therapeutic equivalent to [bisoprolol](#) for

Essential hypertension

Antiarrhythmic medicines

Therapeutic equivalent to [bisoprolol](#) for

Cardiac arrhythmia

Medicines used in heart failure

Therapeutic equivalent to [bisoprolol](#) for

Heart failure

Chlorothiazide

Section

Antihypertensive medicines

Therapeutic equivalent to [hydrochlorothiazide](#) for

Essential hypertension

Medicines used in heart failure

Therapeutic equivalent to [hydrochlorothiazide](#) for

Heart failure

Chlortalidone

Section

Antihypertensive medicines

Medicines used in heart failure

Indications

Therapeutic equivalent to [hydrochlorothiazide](#) for

Essential hypertension

Therapeutic equivalent to [hydrochlorothiazide](#) for

Heart failure

Clopidogrel

Section

Anti-platelet medicines

- Oral > Solid: 75 mg; 300 mg

Indications

Acute ischaemic heart disease

Presence of coronary angioplasty implant or graft

Digitoxin

Section

Medicines used in heart failure

- Parenteral > General injections > IV: 0.2 mg in 1 mL ampoule
- Oral > Solid: 50 µg; 100 µg

Indications

Heart failure

Digoxin

Section

Antiarrhythmic medicines

- Parenteral > General injections > IV: 250 µg per mL in 2 mL ampoule
- Oral > Liquid: 50 µg per mL
- Oral > Solid: 62.5 µg; 250 µg

Medicines used in heart failure

- Parenteral > General injections > IV: 250 µg per mL in 2 mL ampoule; 100 µg per mL in 1 mL ampoule
- Oral > Liquid: 50 µg per mL
- Oral > Solid > tablet: 62.5 µg; 250 µg; 125 µg

Indications

Cardiac arrhythmia

Heart failure

Dopamine

Section

Medicines used in heart failure

- Parenteral > General injections > IV: 40 mg per mL in 5 mL vial (hydrochloride)

Indications

Cardiogenic shock

Heart failure

Enalapril

Section	Indications
Antihypertensive medicines ◦ Oral > Liquid: 1 mg per mL (as hydrogen maleate) ◦ Oral > Solid > tablet: 2.5 mg (as hydrogen maleate); 5 mg (as hydrogen maleate); 10 mg (as hydrogen maleate)	Essential hypertension
Medicines used in heart failure ◦ Oral > Solid > tablet: 2.5 mg (as hydrogen maleate); 5 mg (as hydrogen maleate); 10 mg (as hydrogen maleate)	Heart failure

Epinephrine

Section	Indications
Antiarrhythmic medicines ◦ Parenteral > General injections > IV: 100 µg per mL in 10 mL ampoule (as acid tartrate or hydrochloride)	Cardiac arrest

Fluvastatin

Section	Indications
Lipid-lowering agents	Therapeutic equivalent to simvastatin for Mixed hyperlipidaemia Therapeutic equivalent to simvastatin for Coronary atherosclerosis

Furosemide

Section	Indications
Medicines used in heart failure ◦ Parenteral > General injections > IV: 10 mg per mL in 2 mL ampoule; 10 mg per mL in 5 mL ampoule ◦ Oral > Liquid: 20 mg per 5 mL; 50 mg per 5 mL ◦ Oral > Solid > tablet: 40 mg; 20 mg	Heart failure

Glyceryl trinitrate

Section	Indications
Antianginal medicines ◦ Oral > Solid: 500 µg sublingual tablet	Angina pectoris

Hydralazine

Section

Antihypertensive medicines

- Parenteral > General injections > IV: 20 mg in ampoule powder for injection (hydrochloride)
- Oral > Solid: 50 mg tablet; 25 mg tablet (hydrochloride)
- Oral > Solid: 50 mg (hydrochloride); 25 mg (hydrochloride)

Indications

Essential hypertension

Gestational hypertension

Hydrochlorothiazide

Section

Antihypertensive medicines

- Oral > Liquid: 50 mg per 5 mL
- Oral > Solid: 25 mg; 12.5 mg

Medicines used in heart failure

- Oral > Liquid: 50 mg per 5 mL
- Oral > Solid: 25 mg

Indications

Essential hypertension

Heart failure

Indapamide

Section

Antihypertensive medicines

Therapeutic equivalent to [hydrochlorothiazide](#) for

Essential hypertension

Medicines used in heart failure

Therapeutic equivalent to [hydrochlorothiazide](#) for

Heart failure

Isoprenaline

Section

Antiarrhythmic medicines

- Parenteral > General injections > IV: 20 µg per mL in ampoule (hydrochloride)
- Oral > Solid: 10 mg (hydrochloride or sulfate); 15 mg (hydrochloride or sulfate)

Indications

Bradycardia

Cardiogenic shock

Isosorbide dinitrate

Section

Antianginal medicines

- Oral > Solid: 5 mg sublingual tablet

Indications

Angina pectoris

Lidocaine

Section

Antiarrhythmic medicines

- Parenteral > General injections > IV: 20 mg per mL in 5 mL ampoule (hydrochloride)

Indications

Ventricular tachyarrhythmia

Lisinopril + amlodipine

Section

Antihypertensive medicines

- Oral > Solid: 10 mg + 5 mg; 20 mg + 5 mg; 20 mg + 10 mg

Indications

Essential hypertension

Lisinopril + hydrochlorothiazide

Section

Antihypertensive medicines

- Oral > Solid: 10 mg + 12.5 mg; 20 mg + 12.5 mg; 20 mg + 25 mg

Indications

Essential hypertension

Losartan

Section

Antihypertensive medicines

- Oral > Solid: 25 mg tablet; 50 mg tablet; 100 mg tablet

Medicines used in heart failure

- Oral > Solid: 25 mg tablet; 50 mg tablet; 100 mg tablet

Indications

Essential hypertension

Heart failure

Lovastatin

Section

Lipid-lowering agents

Indications

Therapeutic equivalent to [simvastatin](#) for

Mixed hyperlipidaemia

Therapeutic equivalent to [simvastatin](#) for

Coronary atherosclerosis

Methyldopa

Section

Antihypertensive medicines

- Oral > Solid: 250 mg

Indications

Essential hypertension

Gestational hypertension

Metoprolol

Section

Antianginal medicines

Antihypertensive medicines

Antiarrhythmic medicines

Medicines used in heart failure

Indications

Therapeutic equivalent to [bisoprolol](#) for

Angina pectoris

Therapeutic equivalent to [bisoprolol](#) for

Essential hypertension

Therapeutic equivalent to [bisoprolol](#) for

Cardiac arrhythmia

Therapeutic equivalent to [bisoprolol](#) for

Heart failure

Nifedipine

Section

Antianginal medicines

- Oral > Solid: 10 mg

Antihypertensive medicines

- Oral > Solid: 10 mg tablet; sustained-release formulations

Indications

Angina pectoris

Essential hypertension

Perindopril + amlodipine + indapamide

Section

Antihypertensive medicines

- Oral > Solid > dosage form: 5 mg (arginine) + 5 mg + 1.25 mg; 5 mg (arginine) + 10 mg + 1.25 mg; 10 mg (arginine) + 5 mg + 2.5 mg; 10 mg (arginine) + 10 mg + 2.5 mg

Indications

Essential hypertension

Pravastatin

Section

Lipid-lowering agents

Indications

Therapeutic equivalent to [simvastatin](#) for

Mixed hyperlipidaemia

Therapeutic equivalent to [simvastatin](#) for

Coronary atherosclerosis

Procainamide

Section

Antiarrhythmic medicines

- Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule (hydrochloride)
- Oral > Solid: 500 mg (hydrochloride); 250 mg (hydrochloride)

Indications

Cardiac arrhythmia

Propranolol

Section

Antianginal medicines

- Parenteral > General injections > IV: 1 mg in 1 mL ampoule (hydrochloride)
- Oral > Solid: 10 mg tablet (hydrochloride); 40 mg tablet (hydrochloride)

Antihypertensive medicines

- Oral > Solid: 80 mg tablet (hydrochloride); 40 mg tablet (hydrochloride)

Antiarrhythmic medicines

- Parenteral > General injections > IV: 1 mg in 1 mL ampoule (hydrochloride)
- Oral > Solid: 10 mg tablet (hydrochloride); 40 mg tablet (hydrochloride)

Indications

Angina pectoris

Essential hypertension

Cardiac arrhythmia

Quinidine

Section

Antiarrhythmic medicines

- Oral > Solid: 200 mg (sulfate)

Indications

Cardiac arrhythmia

Simvastatin

Section

Lipid-lowering agents

- Oral > Solid: 5 mg; 10 mg; 20 mg; 40 mg

Indications

Mixed hyperlipidaemia

Coronary atherosclerosis

Sodium nitroprusside

Section

Antihypertensive medicines

- Parenteral > General injections > IV: 50 mg in ampoule powder for infusion

Indications

Hypertensive crisis

Spironolactone

Section

Medicines used in heart failure

- Oral > Solid > tablet: 25 mg

Indications

Heart failure

Streptokinase

Section

Thrombolytic medicines

- Parenteral > General injections > IV: 1.5 million IU in vial powder for injection

Indications

Acute myocardial infarction

Telmisartan + amlodipine

Section

Antihypertensive medicines

- Oral > Solid: 40 mg + 5 mg; 80 mg + 5 mg; 80 mg + 10 mg

Indications

Essential hypertension

Telmisartan + hydrochlorothiazide

Section

Antihypertensive medicines

- Oral > Solid: 40 mg + 12.5 mg; 80 mg + 12.5 mg; 80 mg + 25 mg

Indications

Essential hypertension

Ticagrelor

Section

Anti-platelet medicines

- Oral > Solid > tablet: 60 mg; 90 mg

Indications

Acute ischaemic heart disease

Presence of coronary angioplasty implant or graft

Torsemide

Section

Medicines used in heart failure

Indications

Therapeutic equivalent to furosemide for

Heart failure

Valsartan + amlodipine + hydrochlorothiazide

Section

Antihypertensive medicines

- Oral > Solid > dosage form: 160 mg + 5 mg + 12.5 mg; 160 mg + 5 mg + 25 mg; 160 mg + 10 mg + 12.5 mg; 160 mg + 10 mg + 25 mg; 320 mg + 10 mg + 25 mg

Indications

Essential hypertension

Verapamil

Section

Antianginal medicines

- Oral > Solid: 40 mg (hydrochloride); 80 mg (hydrochloride)

Antiarrhythmic medicines

- Parenteral > General injections > IV: 2.5 mg per mL in 2 mL ampoule (hydrochloride)
- Oral > Solid: 40 mg (hydrochloride); 80 mg (hydrochloride)

Indications

Angina pectoris

Supraventricular tachyarrhythmia