Model List of Essential Medicines

Amidotrizoate

Section	Indications
Diagnostic agents > Radiocontrast media ◦ Parenteral > General injections > unspecified: 140 to 420 mg iodine per mL in 20 mL ampoule (as sodium or meglumine salt)	Amidotrizoate

Atropine

Section	Indications
Diagnostic agents > Ophthalmic medicines	Therapeutic equivalent to tropicamide for Cycloplegic drug

Barium sulfate

Section	Indications
Diagnostic agents > Radiocontrast media ◦ Oral > Liquid: Aqueous suspension	Barium sulfate with suspending agents

Cyclopentolate

Section	Indications
Diagnostic agents > Ophthalmic medicines	Therapeutic equivalent to tropicamide for Cycloplegic drug

Fluorescein

Section	Indications
Diagnostic agents > Ophthalmic medicines ◦ Local > Ophthalmological > Solution (eye drops): 1% (sodium salt)	Fluorescein

Iohexol

Section	Indications
Diagnostic agents > Radiocontrast media ◦ Parenteral > General injections > unspecified: 140 to 350 mg iodine per mL in 5 mL ampoule; 140 to 350 mg iodine per mL in 10 mL ampoule; 140 to 350 mg iodine per mL in 20 mL ampoule	X-ray contrast media, iodinated

Meglumine iotroxate

Section	Indications
Diagnostic agents > Radiocontrast media ◦ Parenteral > General injections > IV: 5 to 8 g iodine (as meglumine) in 100 to 250 mL	X-ray contrast media, iodinated

Tropicamide

Section	Indications
Diagnostic agents > Ophthalmic medicines ◦ Local > Ophthalmological > Solution (eye drops): 0.5%	Cycloplegic drug