



Model List of Essential Medicines



Amidotrizoate [General information](#)

Section

Diagnostic agents > Radiocontrast media

- Parenteral > General injections > unspecified: 140 to 420 mg iodine per mL in 20 mL ampoule (as sodium or meglumine salt)

Indications

[Amidotrizoate](#)



Atropine [General information](#)

Section

Diagnostic agents > Ophthalmic medicines

Indications

Therapeutic equivalent to [tropicamide](#) for [Cycloplegic drug](#)



Barium sulfate [General information](#)

Section

Diagnostic agents > Radiocontrast media

- Oral > Liquid: Aqueous suspension

Indications

[Barium sulfate with suspending agents](#)



Cyclopentolate [General information](#)

Section

Diagnostic agents > Ophthalmic medicines

Indications

Therapeutic equivalent to [tropicamide](#) for [Cycloplegic drug](#)



Fluorescein [General information](#)

Section

Diagnostic agents > Ophthalmic medicines

- Local > Ophthalmological > Solution (eye drops): 1% (sodium salt)

Indications

[Fluorescein](#)



Iohexol [General information](#)

Section

Diagnostic agents > Radiocontrast media

- Parenteral > General injections > unspecified: 140 to 350 mg iodine per mL in 5 mL ampoule; 140 to 350 mg iodine per mL in 10 mL ampoule; 140 to 350 mg iodine per mL in 20 mL ampoule

Indications

[X-ray contrast media, iodinated](#)



Meglumine iotroxate [General information](#)

Section

Diagnostic agents > Radiocontrast media

- Parenteral > General injections > IV: 5 to 8 g iodine (as meglumine) in 100 to 250 mL

Indications

[X-ray contrast media, iodinated](#)



Tropicamide [General information](#)

Section

Diagnostic agents > Ophthalmic medicines

- Local > Ophthalmological > Solution (eye drops): 0.5%

Indications

[Cycloplegic drug](#)