

Alcuronium**Section**

Muscle relaxants (peripherally-acting) and cholinesterase inhibitors

- Parenteral > General injections > IV: 5 mg per mL in 2 mL ampoule (alcuronium chloride)

Indications

Muscle relaxants

Atracurium**Section**

Muscle relaxants (peripherally-acting) and cholinesterase inhibitors

- Parenteral > General injections > IV: 10 mg per mL (besylate)

IndicationsTherapeutic equivalent to [vecuronium](#) for

Muscle relaxants

Muscle relaxants

Gallamine**Section**

Muscle relaxants (peripherally-acting) and cholinesterase inhibitors

- Parenteral > General injections > IV: 40 mg per mL in 2 mL ampoule (gallamine triethiodide)

Indications

Muscle relaxants

Neostigmine**Section**

Muscle relaxants (peripherally-acting) and cholinesterase inhibitors

- Parenteral > General injections > unspecified: 0.5 mg per mL in 1 mL ampoule (neostigmine metilsulfate); 2.5 mg per mL in 1 mL ampoule (neostigmine metilsulfate)
- Oral > Solid > tablet: 15 mg (neostigmine bromide)

Indications

Anticholinesterase reversible

Myasthenia gravis

Pyridostigmine**Section**

Muscle relaxants (peripherally-acting) and cholinesterase inhibitors

- Parenteral > General injections > IV: 5 mg per mL in ampoule or vial (pyridostigmine bromide)
- Oral > Solid > tablet: 60 mg scored (pyridostigmine bromide)

Indications

Myasthenia gravis

Suxamethonium

Section

Muscle relaxants (peripherally-acting) and cholinesterase inhibitors

- Parenteral > General injections > IV: 50 mg per mL in 2 mL ampoule (chloride)

Indications

Muscle relaxants

Tubocurarine

Section

Muscle relaxants (peripherally-acting) and cholinesterase inhibitors

- Parenteral > General injections > IV: 10 mg per mL in 1.5 mL ampoule (tubocurarine chloride)

Indications

Muscle relaxants

Vecuronium

Section

Muscle relaxants (peripherally-acting) and cholinesterase inhibitors

- Parenteral > General injections > IV: 10 mg in vial powder for injection (vecuronium bromide)

Indications

Muscle relaxants