



Model List of Essential Medicines



Acamprosate calcium [General information](#)

Section

Medicines for alcohol use disorders

- Oral > Solid > tablet: 333 mg

Indications

[Disorders due to use of alcohol](#)



Buprenorphine [General information](#)

Section

Medicines for opioid use disorders

Indications

Therapeutic equivalent to [methadone](#) for [Opioid dependence](#)



Bupropion [General information](#)

Section

Medicines for nicotine use disorders

- Oral > Solid: 150 mg sustained-release (hydrochloride)

Indications

[Nicotine dependence](#)



Cytisine (cytisinicline) [General information](#)

Section

Medicines for nicotine use disorders

- Oral > Solid > tablet: 1.5 mg

Indications

[Nicotine dependence](#)



Methadone [General information](#)

Section

Medicines for opioid use disorders

- Oral > Liquid: 5 mg per 5 mL (methadone hydrochloride); 10 mg per 5 mL (methadone hydrochloride); 5 mg per mL concentrate for oral liquid (methadone hydrochloride); 10 mg per mL concentrate for oral liquid (methadone hydrochloride)

Indications

[Opioid dependence](#)



Naltrexone [General information](#)

Section

Medicines for alcohol use disorders

- Parenteral > General injections > IM: 380 mg in vial (extended-release)
- Oral > Solid > tablet: 50 mg

Indications

[Disorders due to use of alcohol](#)



Nicotine replacement therapy [General information](#)

Section

Medicines for nicotine use disorders

- Local > Topical > Transdermal patch: 5 to 30 mg per 16 hour; 7 to 21 mg per 24 hour
- Oral > Solid > lozenge: 2 mg; 4 mg lozenge
- Local > Buccal > Oral spray: 1 mg per actuation
- Local > Buccal > Chewing gum: 2 mg; 4 mg

Indications

[Nicotine dependence](#)



Varenicline [General information](#)

Section

Medicines for nicotine use disorders

- Oral > Solid: 0.5 mg; 1 mg

Indications

[Nicotine dependence](#)