



Model List of Essential Medicines



Acetylsalicylic acid [General information](#)

Section

Medicines for juvenile joint diseases

- Local > Rectal > Suppository: 150 mg; 300 mg
- Oral > Solid > dispersible tablet: 75 mg; 300 mg; 500 mg
- Oral > Solid > tablet: 75 to 500 mg

Indications

[Acute rheumatic fever without mention of heart involvement](#) [Mucocutaneous lymph node syndrome](#) [Juvenile idiopathic arthritis](#)



Adalimumab [General information](#)

Section

Medicines for juvenile joint diseases

- Parenteral > General injections > SC: 40 mg per 0.8 mL; 40 mg per 0.4 mL; 10 mg per 0.2 mL; 20 mg per 0.4 mL

Indications

[Juvenile idiopathic arthritis](#)



Allopurinol [General information](#)

Section

Medicines used to treat gout

- Oral > Solid: 100 mg

Indications

[Gout](#)



Anakinra [General information](#)

Section

Medicines for juvenile joint diseases

- Parenteral > General injections > SC: 100 mg per 0.67 mL in pre-filled syringe

Indications

[Juvenile systemic arthritis](#)



Azathioprine [General information](#)

Section

Disease-modifying anti-rheumatic drugs (DMARDs)

- Oral > Solid > tablet: 50 mg (scored)

Indications

[Rheumatoid arthritis, serology unspecified](#) [Psoriatic arthritis](#)

Certolizumab pegol [General information](#)

Section

Medicines for juvenile joint diseases

Indications

Therapeutic equivalent to [adalimumab](#) for [Juvenile idiopathic arthritis](#)

Chloroquine [General information](#)

Section

Disease-modifying anti-rheumatic drugs (DMARDs)

- Oral > Solid: 100 mg tablet (as phosphate or sulfate); 150 mg tablet (as phosphate or sulfate)

Indications

[Rheumatoid arthritis](#)

Colchicine [General information](#)

Section

Medicines used to treat gout

- Oral > Solid: 0.5 mg

Indications

[Gout](#)

Etanercept [General information](#)

Section

Medicines for juvenile joint diseases

Indications

Therapeutic equivalent to [adalimumab](#) for [Juvenile idiopathic arthritis](#)

Golimumab [General information](#)

Section

Medicines for juvenile joint diseases

Indications

Therapeutic equivalent to [adalimumab](#) for [Juvenile idiopathic arthritis](#)

Hydroxychloroquine [General information](#)

Section

Disease-modifying anti-rheumatic drugs (DMARDs)

- Oral > Solid: 200 mg (as sulfate)

Indications

[Systemic lupus erythematosus](#) [Cutaneous lupus erythematosus of unspecified type](#)

Infliximab [General information](#)

Section

Medicines for juvenile joint diseases

Indications

Therapeutic equivalent to [adalimumab](#) for [Juvenile idiopathic arthritis](#)

Methotrexate [General information](#)

Section

Disease-modifying anti-rheumatic drugs (DMARDs)

- Oral > Solid > tablet: 2.5 mg (as sodium salt)

Indications

[Rheumatoid arthritis, serology unspecified](#)

Section

Medicines for juvenile joint diseases

- Oral > Solid > tablet: 2.5 mg (as sodium salt)

Indications

[Juvenile idiopathic arthritis](#)

Penicillamine [General information](#)

Section

Disease-modifying anti-rheumatic drugs (DMARDs)

- Oral > Solid: 250 mg

Indications

[Rheumatoid arthritis, serology unspecified](#)

Probenecid [General information](#)

Section

Medicines used to treat gout

- Oral > Solid: 500 mg

Indications

[Gout](#)

Sulfasalazine [General information](#)

Section

Disease-modifying anti-rheumatic drugs (DMARDs)

- Oral > Solid: 500 mg

Indications

[Rheumatoid arthritis, serology unspecified](#)

Tocilizumab [General information](#)

Section

Medicines for juvenile joint diseases

- Parenteral > General injections > IV: 80 mg per 4 mL in vial; 200 mg per 10 mL in vial; 400 mg per 20 mL in vial
- Parenteral > General injections > SC: 162 mg per 0.9 mL in pre-filled syringe

Indications

[Juvenile systemic arthritis](#)

Triamcinolone hexacetonide [General information](#)

Section

Medicines for juvenile joint diseases

- Parenteral > Locoregional injections > Intra-articular: 20 mg per mL in vial

Indications

[Juvenile idiopathic arthritis](#)