

Abacavir + dolutegravir + lamivudine

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid > tablet: 60 mg (as sulfate) + 5 mg + 30 mg (dispersible, scored)	Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Abacavir + lamivudine

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 120 mg (as sulfate) + 60 mg tablet (dispersible, scored)	Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Abacavir + lamivudine + lopinavir + ritonavir

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 30 mg + 15 mg + 40 mg + 10 mg capsule containing oral granules	Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Cobicistat + elvitegravir + emtricitabine + tenofovir disoproxil fumarate

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 150 mg + 150 mg + 200 mg + 300 mg (tenofovir disoproxil fumarate equivalent to 245 mg tenofovir disoproxil)	Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Cobicistat + elvitegravir + emtricitabine+ tenofovir alafenamide

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 150 mg + 150 mg + 200 mg + 10 mg	Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Dolutegravir + lamivudine + tenofovir

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 50 mg + 300 mg + 300 mg (tenofovir disoproxil fumarate equivalent to 245 mg tenofovir disoproxil)	Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Efavirenz + emtricitabine + tenofovir

Section

Fixed-dose combinations of antiretrovirals

- Oral > Solid: 600 mg + 200 mg + 300 mg (tenofovir disoproxil fumarate equivalent to 245 mg tenofovir disoproxil)

Indications

Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Efavirenz + lamivudine + tenofovir

Section

Fixed-dose combinations of antiretrovirals

- Oral > Solid: 400 mg + 300 mg + 300 mg (tenofovir disoproxil fumarate, equivalent to 245 mg tenofovir disoproxil)

Indications

Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Therapeutic equivalent to efavirenz + emtricitabine + tenofovir for

Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Emtricitabine + rilpivirine + tenofovir alafenamide

Section

Fixed-dose combinations of antiretrovirals

- Oral > Solid: 200 mg + 25 mg + 25 mg

Indications

Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Emtricitabine + rilpivirine + tenofovir disoproxil fumarate

Section

Fixed-dose combinations of antiretrovirals

- Oral > Solid: 200 mg + 25 mg + 300 mg (tenofovir disoproxil fumarate equivalent to 245 mg tenofovir disoproxil)

Indications

Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Emtricitabine + tenofovir

Section

Fixed-dose combinations of antiretrovirals

- Oral > Solid: 200 mg + 300 mg (tenofovir disoproxil fumarate equivalent to 245 mg tenofovir disoproxil)

Indications

Contact with or exposure to human immunodeficiency virus

Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Emtricitabine + tenofovir alafenamide

Section

Fixed-dose combinations of antiretrovirals

- Oral > Solid: 200 mg + 10 mg; 200 mg + 25 mg

Indications

Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Lamivudine + nevirapine + zidovudine

Section

Fixed-dose combinations of antiretrovirals

- Oral > Solid: 30 mg + 50 mg + 60 mg Tablet; 150 mg + 200 mg + 300 mg Tablet

Indications

Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Lamivudine + tenofovir

Section

Fixed-dose combinations of antiretrovirals

Indications

Therapeutic equivalent to emtricitabine + tenofovir for

Contact with or exposure to human immunodeficiency virus

Therapeutic equivalent to emtricitabine + tenofovir for

Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Lamivudine + zidovudine

Section

Fixed-dose combinations of antiretrovirals

- Oral > Solid: 30 mg + 60 mg Tablet; 150 mg + 300 mg Tablet

Indications

Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified