Model List of Essential Medicines

Atezolizumab

Section	Indications
Immunomodulators ◦ Parenteral > General injections > IV: 840 mg per 14 mL in vial concentrate for solution for infusion; 1200 mg per 20 mL in vial concentrate for solution for infusion ◦ Parenteral > General injections > IV: 1.2 g per 20 mL concentrate solution for infusion	Therapeutic equivalent to pembrolizumab for Adenocarcinoma of stomach Hepatocellular carcinoma of liver Therapeutic equivalent to pembrolizumab for Other specified malignant neoplasms of bronchus or lung Other specified malignant neoplasms of bronchus or lung

Bevacizumab

Section	Indications
Immunomodulators ◦ Parenteral > General injections > IV: 25 mg per mL in 4 mL vial concentrate for solution for infusion; 25 mg per mL in 16 mL vial concentrate for solution for infusion	Hepatocellular carcinoma of liver

Blinatumomab

Section	Indications
Immunomodulators ◦ Parenteral > General injections > IV: 35 µg in vial; 38.5 µg in vial	B lymphoblastic leukaemia or lymphoma, not elsewhere classified

Cemiplimab

Section	Indications
Immunomodulators ◦ Parenteral > General injections > IV: 50 mg per mL in 7 mL vial concentrate for solution for infusion	Therapeutic equivalent to pembrolizumab for Adenocarcinoma of stomach Therapeutic equivalent to pembrolizumab for Other specified malignant neoplasms of bronchus or lung Other specified malignant neoplasms of bronchus or lung

Daratumumab

Section	Indications
Immunomodulators ◦ Parenteral > General injections > IV: 100 mg per 5 mL; 400 mg per 20 mL	Plasma cell myeloma

Dostarlimab

Section	Indications
Immunomodulators ◦ Parenteral > General injections > IV: 50 mg per mL in 10 mL vial concentrate for solution for infusion	Other specified malignant neoplasms of corpus uteri

Durvalumab

Section	Indications
Immunomodulators ◦ Parenteral > General injections > IV: 50 mg per mL in 2.4 mL vial concentrate solution for infusion; 50 mg per mL in 10 mL vial concentrate solution for infusion	Hepatocellular carcinoma of liver Malignant neoplasms of biliary tract, distal bile duct Malignant neoplasm of intrahepatic bile ducts Malignant neoplasms of gallbladder Malignant neoplasms of proximal biliary tract, cystic duct Other specified malignant neoplasms of bronchus or lung

Filgrastim

Section	Indications
Immunomodulators ◦ Parenteral > General injections > SC: 120 µg per 0.2 mL in prefilled syringe; 300 µg per 0.5 mL in prefilled syringe; 480 µg per 0.8 mL in prefilled syringe; 300 µg per mL in 1 mL vial; 480 µg per 1.6 mL in 1.6 mL vial	Acquired neutropaenia

Ipilimumab

Section	Indications
Immunomodulators ◦ Parenteral > General injections > IV: 5 mg per mL in 10 mL vial concentrate for solution for infusion; 20 mg per mL in 40 mL vial concentrate for solution for infusion	Renal cell carcinoma of kidney, except renal pelvis Squamous cell carcinoma of oesophagus Melanoma of skin Other specified malignant neoplasms of bronchus or lung

Lenalidomide

Section	Indications
Immunomodulators ◦ Oral > Solid: 25 mg	Plasma cell myeloma

Nivolumab

Section	Indications
Immunomodulators ◦ Parenteral > General injections > IV: 10 mg per mL concentrate solution for infusion ◦ Parenteral > General injections > IV: 10 mg per mL in 4 mL vial concentrate solution for infusion; 10 mg per mL in 10 mL vial concentrate solution for infusion; 10 mg per mL in 12 mL vial concentrate solution for infusion; 10 mg per mL in 24 mL vial concentrate solution for infusion ◦ Parenteral > General injections > IV: 10 mg per mL in 24 mL vial concentrate solution for infusion; 10 mg per mL in 4 mL vial concentrate solution for infusion; 10 mg per mL in 10 mL vial concentrate solution for infusion; 10 mg per mL in 12 mL vial concentrate solution for infusion	Therapeutic equivalent to pembrolizumab for Melanoma of skin Melanoma of skin Renal cell carcinoma of kidney, except renal pelvis Squamous cell carcinoma of oesophagus Adenocarcinoma of oesophagogastric junction Adenocarcinoma of stomach Other specified malignant neoplasms of bronchus or lung

Pegfilgrastim

Section	Indications
Immunomodulators ◦ Parenteral > General injections > SC: 6 mg per 0.6 mL in pre-filled syringe	Acquired neutropaenia

Pembrolizumab

Section	Indications
Immunomodulators ◦ Parenteral > General injections > IV: 25 mg per mL in 4 mL vial ◦ Parenteral > General injections > IV: 25 mg per mL in 4 mL vial concentrate for solution for infusion ◦ Parenteral > General injections > IV: 25 mg per mL in 4 mL vial powder for injection	Melanoma of skin Other specified malignant neoplasms of large intestine Malignant neoplasms of cervix uteri Adenocarcinoma of stomach Malignant neoplasms of oesophagogastric junction Squamous cell carcinoma of hypopharynx and variants Squamous cell carcinoma of other or ill-defined sites in the lip, oral cavity or pharynx Squamous cell carcinoma of oropharynx Squamous cell carcinoma of nasopharynx Therapeutic equivalent to nivolumab for Melanoma of skin Renal cell carcinoma of kidney, except renal pelvis Carcinoma of breast, specialised type Squamous cell carcinoma of oesophagus Other specified malignant neoplasms of bronchus or lung

Thalidomide

Section	Indications
Immunomodulators ◦ Oral > Solid: 50 mg	Plasma cell myeloma

Tislelizumab

Section	Indications
Immunomodulators ◦ Parenteral > General injections > IV: 100 mg per 10 mL ◦ Parenteral > General injections > IV: 100 mg per 10 mL concentrate for solution for infusion ◦ Parenteral > General injections > IV: 100 mg per 10 mL contentrate for solution for infusion	Malignant neoplasms of urinary tract Squamous cell carcinoma of oesophagus Other specified malignant neoplasms of bronchus or lung Hodgkin lymphoma

Toripalimab

Section	Indications
Immunomodulators ◦ Parenteral > General injections > IV: 240 mg per 6 mL in vial	Squamous cell carcinoma of oesophagus Malignant neoplasms of nasopharynx

Tremelimumab

Section	Indications
Immunomodulators ◦ Parenteral > General injections > IV: 20 mg per mL in 1.25 mL vial concentrate solution for infusion; 20 mg per mL in 15 mL vial concentrate solution for infusion	Hepatocellular carcinoma of liver Other specified malignant neoplasms of bronchus or lung