



Model List of Essential Medicines



Atezolizumab [General information](#)

Section

Immunomodulators

- Parenteral > General injections > IV: 1.2 g per 20 mL concentrate solution for infusion
- Parenteral > General injections > IV: 840 mg per 14 mL in vial concentrate for solution for infusion; 1200 mg per 20 mL in vial concentrate for solution for infusion

Indications

[Other specified malignant neoplasms of bronchus or lung](#) Therapeutic equivalent to [pembrolizumab](#) for [Other specified malignant neoplasms of bronchus or lung](#) Therapeutic equivalent to [pembrolizumab](#) for [Adenocarcinoma of stomach Hepatocellular carcinoma of liver](#)



Bevacizumab [General information](#)

Section

Immunomodulators

- Parenteral > General injections > IV: 25 mg per mL in 4 mL vial concentrate for solution for infusion; 25 mg per mL in 16 mL vial concentrate for solution for infusion

Indications

[Hepatocellular carcinoma of liver](#)



Blinatumomab [General information](#)

Section

Immunomodulators

- Parenteral > General injections > IV: 35 µg in vial; 38.5 µg in vial

Indications

[B lymphoblastic leukaemia or lymphoma, not elsewhere classified](#)



Cemiplimab [General information](#)

Section

Immunomodulators

- Parenteral > General injections > IV: 50 mg per mL in 7 mL vial concentrate for solution for infusion

Indications

Therapeutic equivalent to [pembrolizumab](#) for [Other specified malignant neoplasms of bronchus or lung](#)
Therapeutic equivalent to [pembrolizumab](#) for [Adenocarcinoma of stomach Other specified malignant neoplasms of bronchus or lung](#)



Daratumumab [General information](#)

Section

Immunomodulators

- Parenteral > General injections > IV: 100 mg per 5 mL; 400 mg per 20 mL

Indications

[Plasma cell myeloma](#)



Dostarlimab [General information](#)

Section

Immunomodulators

- Parenteral > General injections > IV: 50 mg per mL in 10 mL vial concentrate for solution for infusion

Indications

[Other specified malignant neoplasms of corpus uteri](#)



Durvalumab [General information](#)

Section

Immunomodulators

- Parenteral > General injections > IV: 50 mg per mL in 2.4 mL vial concentrate solution for infusion; 50 mg per mL in 10 mL vial concentrate solution for infusion

Indications

[Hepatocellular carcinoma of liver Malignant neoplasms of biliary tract, distal bile duct Malignant neoplasm of intrahepatic bile ducts Malignant neoplasms of gallbladder Other specified malignant neoplasms of bronchus or lung Malignant neoplasms of proximal biliary tract, cystic duct](#)



Filgrastim [General information](#)

Section

Immunomodulators

- Parenteral > General injections > SC: 120 µg per 0.2 mL in prefilled syringe; 300 µg per 0.5 mL in prefilled syringe; 480 µg per 0.8 mL in prefilled syringe; 300 µg per mL in 1 mL vial; 480 µg per 1.6 mL in 1.6 mL vial

Indications

[Acquired neutropaenia](#)



Ipilimumab [General information](#)

Section

Immunomodulators

- Parenteral > General injections > IV: 5 mg per mL in 10 mL vial concentrate for solution for infusion; 20 mg per mL in 40 mL vial concentrate for solution for infusion

Indications

[Renal cell carcinoma of kidney, except renal pelvis Squamous cell carcinoma of oesophagus Melanoma of skin Other specified malignant neoplasms of bronchus or lung](#)



Lenalidomide [General information](#)

Section

Immunomodulators

- Oral > Solid: 25 mg

Indications

[Plasma cell myeloma](#)



Nivolumab [General information](#)

Section

Immunomodulators

- Parenteral > General injections > IV: 10 mg per mL concentrate solution for infusion
- Parenteral > General injections > IV: 10 mg per mL in 4 mL vial concentrate solution for infusion; 10 mg per mL in 10 mL vial concentrate solution for infusion; 10 mg per mL in 12 mL vial concentrate solution for infusion; 10 mg per mL in 24 mL vial concentrate solution for infusion
- Parenteral > General injections > IV: 10 mg per mL in 24 mL vial concentrate solution for infusion; 10 mg per mL in 4 mL vial concentrate solution for infusion; 10 mg per mL in 10 mL vial concentrate solution for infusion; 10 mg per mL in 12 mL vial concentrate solution for infusion

Indications

Therapeutic equivalent to [pembrolizumab](#) for [Melanoma of skin Melanoma of skin Renal cell carcinoma of kidney, except renal pelvis Other specified malignant neoplasms of bronchus or lung Squamous cell carcinoma of oesophagus Adenocarcinoma of oesophagogastric junction Adenocarcinoma of stomach](#)



Pegfilgrastim [General information](#)

Section

Immunomodulators

- Parenteral > General injections > SC: 6 mg per 0.6 mL in pre-filled syringe

Indications

[Acquired neutropaenia](#)



Pembrolizumab [General information](#)

Section

Immunomodulators

- Parenteral > General injections > IV: 25 mg per mL in 4 mL vial
- Parenteral > General injections > IV: 25 mg per mL in 4 mL vial powder for injection
- Parenteral > General injections > IV: 25 mg per mL in 4 mL vial concentrate for solution for infusion

Indications

[Melanoma of skin](#) [Other specified malignant neoplasms of large intestine](#) [Malignant neoplasms of cervix uteri](#) [Other specified malignant neoplasms of bronchus or lung](#) [Adenocarcinoma of stomach](#) [Malignant neoplasms of oesophagogastric junction](#) [Squamous cell carcinoma of hypopharynx and variants](#) [Squamous cell carcinoma of other or ill-defined sites in the lip, oral cavity or pharynx](#) [Squamous cell carcinoma of oropharynx](#) [Squamous cell carcinoma of nasopharynx](#) Therapeutic equivalent to [nivolumab](#) for [Melanoma of skin](#) [Renal cell carcinoma of kidney, except renal pelvis](#) [Carcinoma of breast, specialised type](#) [Squamous cell carcinoma of oesophagus](#)



Thalidomide [General information](#)

Section

Immunomodulators

- Oral > Solid: 50 mg

Indications

[Plasma cell myeloma](#)



Tislelizumab [General information](#)

Section

Immunomodulators

- Parenteral > General injections > IV: 100 mg per 10 mL
- Parenteral > General injections > IV: 100 mg per 10 mL concentrate for solution for infusion
- Parenteral > General injections > IV: 100 mg per 10 mL concentrate for solution for infusion

Indications

[Malignant neoplasms of urinary tract](#) [Hodgkin lymphoma](#) [Other specified malignant neoplasms of bronchus or lung](#) [Squamous cell carcinoma of oesophagus](#)



Toripalimab [General information](#)

Section

Immunomodulators

- Parenteral > General injections > IV: 240 mg per 6 mL in vial

Indications

[Malignant neoplasms of nasopharynx](#) [Squamous cell carcinoma of oesophagus](#)



Tremelimumab [General information](#)

Section

Immunomodulators

- Parenteral > General injections > IV: 20 mg per mL in 1.25 mL vial concentrate solution for infusion; 20 mg per mL in 15 mL vial concentrate solution for infusion

Indications

[Hepatocellular carcinoma of liver](#) [Other specified malignant neoplasms of bronchus or lung](#)