Liste modèle des médicaments essentiels

Beinaglutide

Section	Indications
Hypoglycaemic agents	Therapeutic equivalent to semaglutide pour Obesity

Canagliflozin

Section	Indications
Hypoglycaemic agents	Therapeutic equivalent to empagliflozin pour Type 2 diabetes mellitus

Dapagliflozin

Section	Indications
Hypoglycaemic agents	Therapeutic equivalent to empagliflozin pour Type 2 diabetes mellitus

Dulaglutide

Section	Indications
Hypoglycaemic agents	Therapeutic equivalent to semaglutide pour Obesity Therapeutic equivalent to semaglutide pour Type 2 diabetes mellitus

Empagliflozin

Section	Indications
Hypoglycaemic agents ◦ Oral > Solid: 10 mg; 25 mg	Type 2 diabetes mellitus

Exenatide

Section	Indications
Hypoglycaemic agents	Therapeutic equivalent to semaglutide pour Obesity

Glibenclamide

Section	Indications
Hypoglycaemic agents ◦ Oral > Solid: 5 mg; 2.5 mg	Type 2 diabetes mellitus

Gliclazide

Section	Indications
Hypoglycaemic agents ◦ Oral > Solid: 30 mg controlled-release; 60 mg controlled-release; 80 mg controlled-release	Type 2 diabetes mellitus

Liraglutide

Section	Indications
Hypoglycaemic agents	Therapeutic equivalent to semaglutide pour Obesity Therapeutic equivalent to semaglutide pour Type 2 diabetes mellitus

Metformin

Section	Indications
Hypoglycaemic agents ◦ Oral > Solid: 500 mg (hydrochloride)	Type 2 diabetes mellitus

Semaglutide

Section	Indications
Hypoglycaemic agents ◦ Parenteral > General injections > SC: 0.25 mg in pre-filled pen injection solution; 0.5 mg in pre-filled pen injection solution; 1 mg in pre-filled pen injection solution; 2 mg in pre-filled pen injection solution; 4 mg in pre-filled pen injection solution ◦ Oral > Solid > tablet: 3 mg ◦ Parenteral > General injections > SC: 0.68 mg per mL; 1.34 mg per mL; 2.68 mg per mL	Obesity Type 2 diabetes mellitus

Tirzepatide

Section	Indications
Hypoglycaemic agents	Therapeutic equivalent to semaglutide pour Obesity Therapeutic equivalent to semaglutide pour Type 2 diabetes mellitus

Tolbutamide

Section	Indications
Hypoglycaemic agents ◦ Oral > Solid: 500 mg	Type 2 diabetes mellitus