

Alprostadil (prostaglandin E1)**Section**

Medicines administered to the neonate [c]

- Parenteral > General injections > IV: 0.5 mg per mL in alcohol

Indications

Other specified congenital anomaly of great arteries including arterial duct

Beractant**Section**

Medicines administered to the neonate [c]

- Respiratory > Suspension: 25 mg per mL for intratracheal instillation

Indications

Respiratory distress syndrome of newborn

Caffeine citrate**Section**

Medicines administered to the neonate [c]

- Parenteral > General injections > IV: 20 mg per mL (equivalent to 10 mg caffeine base per mL)
- Oral > Liquid: 20 mg per mL (equivalent to 10 mg caffeine base per mL)

Indications

Apnoea of newborn

Chlorhexidine**Section**

Medicines administered to the neonate [c]

- Local > Topical > Solution: 7.1% chlorhexidine digluconate (delivering 4% chlorhexidine)
- Local > Topical > Gel: 7.1% chlorhexidine digluconate (delivering 4% chlorhexidine)

Indications

Omphalitis of newborn

Dinoprostone (prostaglandin E2)**Section**

Medicines administered to the neonate [c]

- Parenteral > General injections > IV: 1 mg per mL

Indications

Therapeutic equivalent to [alprostadil \(prostaglandin E1\)](#) for

Other specified congenital anomaly of great arteries including arterial duct

Other specified congenital anomaly of great arteries including arterial duct

Ibuprofen

Section

Medicines administered to the neonate [c]

- Parenteral > General injections > IV: 5 mg per mL

Indications

Patent arterial duct

Indometacin

Section

Medicines administered to the neonate [c]

Indications

Therapeutic equivalent to [ibuprofen](#) for

Patent arterial duct

Porcatant alfa

Section

Medicines administered to the neonate [c]

- Respiratory > Suspension: 80 mg per mL for intratracheal instillation

Indications

Respiratory distress syndrome of newborn