Model List of Essential Medicines

Alprostadil (prostaglandin E1)

Section	Indications
Medicines administered to the neonate [c] ◦ Parenteral > General injections > IV: 0.5 mg per mL in alcohol	Other specified congenital anomaly of great arteries including arterial duct

Beractant

Section	Indications
Medicines administered to the neonate [c] ◦ Respiratory > Suspension: 25 mg per mL for intratracheal instillation	Respiratory distress syndrome of newborn

Caffeine citrate

Section	Indications
Medicines administered to the neonate [c] ◦ Parenteral > General injections > IV: 20 mg per mL (equivalent to 10 mg caffeine base per mL) ◦ Oral > Liquid: 20 mg per mL (equivalent to 10 mg caffeine base per mL)	Apnoea of newborn

Chlorhexidine

Section	Indications
Medicines administered to the neonate [c] ◦ Local > Topical > Solution: 7.1% chlorhexidine digluconate (delivering 4% chlorhexidine) ◦ Local > Topical > Gel: 7.1% chlorhexidine digluconate (delivering 4% chlorhexidine)	Omphalitis of newborn

Dinoprostone (prostaglandin E2)

Section	Indications
Medicines administered to the neonate [c] ◦ Parenteral > General injections > IV: 1 mg per mL	Therapeutic equivalent to alprostadil (prostaglandin E1) for Other specified congenital anomaly of great arteries including arterial duct Other specified congenital anomaly of great arteries including arterial duct

Ibuprofen

Section	Indications
Medicines administered to the neonate [c] ◦ Parenteral > General injections > IV: 5 mg per mL	Patent arterial duct

Indometacin

Section	Indications
Medicines administered to the neonate [c]	Therapeutic equivalent to ibuprofen for Patent arterial duct

Porcatant alfa

Section	Indications
Medicines administered to the neonate [c] ◦ Respiratory > Suspension: 80 mg per mL for intratracheal instillation	Respiratory distress syndrome of newborn