Model List of Essential Medicines

Acenocoumarol

Section	Indications
Medicines affecting coagulation	Therapeutic equivalent to warfarin for Venous thromboembolism Therapeutic equivalent to warfarin for Cerebral ischaemic stroke due to embolic occlusion Therapeutic equivalent to warfarin for Atrial fibrillation Therapeutic equivalent to warfarin for Other specified diseases of arteries or arterioles

Apixaban

Section	Indications
Medicines affecting coagulation	Therapeutic equivalent to dabigatran for Other specified atrial fibrillation Therapeutic equivalent to dabigatran for Venous thromboembolism

Dabigatran

Section	Indications
Medicines affecting coagulation ◦ Oral > Solid: 110 mg; 150 mg	Other specified atrial fibrillation Venous thromboembolism

Dalteparin

Section	Indications
Medicines affecting coagulation	Therapeutic equivalent to enoxaparin for Acute ischaemic heart disease Therapeutic equivalent to enoxaparin for Venous thromboembolism

Desmopressin

Section	Indications
Medicines affecting coagulation ◦ Parenteral > General injections > IV: 4 µg per mL in 1 mL ampoule (acetate) ◦ Local > Nasal > Spray: 150 µg per actuation (acetate)	Haemophilia A Von Willebrand disease

Edoxaban

Section	Indications
Medicines affecting coagulation	Therapeutic equivalent to dabigatran for Other specified atrial fibrillation Therapeutic equivalent to dabigatran for Venous thromboembolism

Emicizumab

Section	Indications
Medicines affecting coagulation ◦ Parenteral > General injections > SC: 12 mg per 0.4 mL in vial; 30 mg per mL in mL vial; 60 mg per 0.4 mL in vial; 105 mg per 0.7 mL in vial; 150 mg per mL in vial; 300 mg per 2 mL in vial	Haemophilia A

Enoxaparin

Section	Indications
Medicines affecting coagulation ◦ Parenteral > General injections > SC: 20 mg per 0.2 mL in prefilled syringe; 40 mg per 0.4 mL in prefilled syringe; 60 mg per 0.6 mL in prefilled syringe; 80 mg per 0.8 mL in prefilled syringe; 100 mg per 1 mL in prefilled syringe; 120 mg per 0.8 mL in prefilled syringe; 150 mg per 1 mL in prefilled syringe; 150 mg per 1 mL in ampoule; 20 mg per 0.2 mL in ampoule; 40 mg per 0.4 mL in ampoule; 60 mg per 0.6 mL in ampoule; 80 mg per 0.8 mL in ampoule; 100 mg per 1 mL in ampoule; 120 mg per 0.8 mL in ampoule	Acute ischaemic heart disease Venous thromboembolism

Heparin sodium

Section	Indications
Medicines affecting coagulation ◦ Parenteral > General injections > IV: 1000 IU per mL in 1 mL ampoule or vial; 20000 IU per mL in 1 mL ampoule or vial; 5000 IU per mL in 1 mL ampoule or vial	Acute ischaemic heart disease Unstable angina Venous thromboembolism

Nadroparin

Section	Indications
Medicines affecting coagulation	Therapeutic equivalent to enoxaparin for Acute ischaemic heart disease Therapeutic equivalent to enoxaparin for Venous thromboembolism

Phytomenadione

Section	Indications
Medicines affecting coagulation ◦ Parenteral > General injections > IV: 10 mg per mL in ampoule; 1 mg per mL in ampoule; 2 mg per 0.2 mL in ampoule mixed micelle solution; 10 mg per mL in ampoule mixed micelle solution; 1 mg per 0.5 mL in ampoule ◦ Oral > Solid > tablet: 5 mg tablet	Haemorrhagic disorder due to circulating anticoagulants or coagulation factors inhibitors Haemorrhagic or haematological disorders of fetus or newborn

Protamine sulfate

Section	Indications
Medicines affecting coagulation ◦ Parenteral > General injections > IV: 10 mg per mL in 5 mL ampoule or vial	Haemorrhagic disorder due to circulating anticoagulants or coagulation factors inhibitors

Rivaroxaban

Section	Indications
Medicines affecting coagulation	Therapeutic equivalent to dabigatran for Other specified atrial fibrillation Therapeutic equivalent to dabigatran for Venous thromboembolism

Tranexamic acid

Section	Indications
Medicines affecting coagulation ◦ Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule	Haemorrhage, not elsewhere classified

Warfarin

Section	Indications
Medicines affecting coagulation ◦ Oral > Solid > tablet: 5 mg (scored) (sodium); 1 mg (scored) (sodium); 2 mg (scored) (sodium); 0.5 mg (scored) (sodium); 3 mg (scored) (sodium) ◦ Oral > Solid > tablet: 5 mg (scored) (sodium); 1 mg (scored) (sodium); 2 mg (scored) (sodium); 0.5 mg (scored) (sodium); 3 mg (scored) (sodium)	Venous thromboembolism Cerebral ischaemic stroke due to embolic occlusion Atrial fibrillation Other specified diseases of arteries or arterioles