Model List of Essential Medicines

Alcuronium

Section	Indications
Muscle relaxants (peripherally-acting) and cholinesterase inhibitors ◦ Parenteral > General injections > IV: 5 mg per mL in 2 mL ampoule (alcuronium chloride)	Muscle relaxants

Atracurium

Section	Indications
Muscle relaxants (peripherally-acting) and cholinesterase inhibitors ◦ Parenteral > General injections > IV: 10 mg per mL (besylate)	Therapeutic equivalent to vecuronium for Muscle relaxants Muscle relaxants

Gallamine

Section	Indications
Muscle relaxants (peripherally-acting) and cholinesterase inhibitors ◦ Parenteral > General injections > IV: 40 mg per mL in 2 mL ampoule (gallamine triethiodide)	Muscle relaxants

Neostigmine

Section	Indications
Muscle relaxants (peripherally-acting) and cholinesterase inhibitors ◦ Parenteral > General injections > unspecified: 0.5 mg per mL in 1 mL ampoule (neostigmine metilsulfate); 2.5 mg per mL in 1 mL ampoule (neostigmine metilsulfate) ◦ Oral > Solid > tablet: 15 mg (neostigmine bromide)	Anticholinesterase reversible Myasthenia gravis

Pyridostigmine

Section	Indications
Muscle relaxants (peripherally-acting) and cholinesterase inhibitors ◦ Parenteral > General injections > IV: 5 mg per mL in ampoule or vial (pyridostigmine bromide) ◦ Oral > Solid > tablet: 60 mg scored (pyridostigmine bromide)	Myasthenia gravis

Suxamethonium

Section	Indications
Muscle relaxants (peripherally-acting) and cholinesterase inhibitors ◦ Parenteral > General injections > IV: 50 mg per mL in 2 mL ampoule (chloride)	Muscle relaxants

Tubocurarine

Section	Indications
Muscle relaxants (peripherally-acting) and cholinesterase inhibitors ◦ Parenteral > General injections > IV: 10 mg per mL in 1.5 mL ampoule (tubocurarine chloride)	Muscle relaxants

Vecuronium

Section	Indications
Muscle relaxants (peripherally-acting) and cholinesterase inhibitors ◦ Parenteral > General injections > IV: 10 mg in vial powder for injection (vecuronium bromide)	Muscle relaxants