# Liste modèle des médicaments essentiels

## Acetazolamide

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Miotics and antiglaucoma medicines<br>◦ Oral > Solid: 250 mg | Primary open-angle glaucoma |

## Aciclovir

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Anti-infective agents<br>◦ Local > Ophthalmological > Ointment: 3% w/w | Herpes simplex keratitis |

## Amikacin

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Anti-infective agents | Therapeutic equivalent to gentamicin pour<br>Other specified conjunctivitis<br>Therapeutic equivalent to gentamicin pour<br>Infectious blepharitis |

## Atropine

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Mydriatics<br>◦ Local > Ophthalmological > Solution (eye drops): 0.1% (sulfate); 0.5% (sulfate); 1% (sulfate) | Anterior uveitis |

## Azithromycin

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Anti-infective agents<br>◦ Local > Ophthalmological > Solution (eye drops): 1.5% eye drops | Trachoma |

## Bevacizumab

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Anti-vascular endothelial growth factor (VEGF) preparations<br>◦ Parenteral > Locoregional injections > Intravitreal: 25 mg per mL | Age-related macular degeneration |

## Carbachol

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Miotics and antiglaucoma medicines | Therapeutic equivalent to pilocarpine pour<br>Acute angle closure with pupillary block<br>Therapeutic equivalent to pilocarpine pour<br>Ocular hypertension<br>Therapeutic equivalent to pilocarpine pour<br>Primary open-angle glaucoma |

## Carmellose

| Section | Indications |
| --- | --- |
| Ophthalmological preparations | Therapeutic equivalent to hypromellose pour<br>Keratoconjunctivitis sicca |

## Chlortetracycline

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Anti-infective agents | Therapeutic equivalent to tetracycline pour<br>Other specified conjunctivitis<br>Therapeutic equivalent to tetracycline pour<br>Infectious keratitis<br>Therapeutic equivalent to tetracycline pour<br>Infectious blepharitis<br>Therapeutic equivalent to tetracycline pour<br>Trachoma |

## Cyclopentolate

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Mydriatics | Therapeutic equivalent to atropine pour<br>Anterior uveitis |

## Epinephrine

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Mydriatics<br>◦ Local > Ophthalmological > Solution (eye drops): 2% (as hydrochloride) | Primary open-angle glaucoma |

## Erythromycin

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Anti-infective agents<br>◦ Local > Ophthalmological > Ointment: 0.5% | Chlamydial conjunctivitis<br>Gonococcal infection of eye |

## Gentamicin

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Anti-infective agents<br>◦ Local > Ophthalmological > Solution (eye drops): 0.3% (gentamicin sulfate) | Other specified conjunctivitis<br>Infectious blepharitis |

## Homatropine

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Mydriatics<br>◦ Local > Ophthalmological > Solution (eye drops): 2% (hydrobromide) | Anterior uveitis<br>Therapeutic equivalent to atropine pour<br>Anterior uveitis |

## Hydrocortisone

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Anti-inflammatory agents<br>◦ Local > Ophthalmological > Ointment: 1% (acetate) | Scleritis<br>Anterior uveitis |

## Hypromellose

| Section | Indications |
| --- | --- |
| Ophthalmological preparations<br>◦ Local > Ophthalmological > Solution (eye drops): 0.3% | Keratoconjunctivitis sicca |

## Idoxuridine

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Anti-infective agents<br>◦ Local > Ophthalmological > Solution (eye drops): 0.1%<br>◦ Local > Ophthalmological > Ointment: 0.2% | Herpes simplex keratitis |

## Kanamycin

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Anti-infective agents | Therapeutic equivalent to gentamicin pour<br>Other specified conjunctivitis<br>Therapeutic equivalent to gentamicin pour<br>Infectious blepharitis |

## Latanoprost

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Miotics and antiglaucoma medicines<br>◦ Local > Ophthalmological > Solution (eye drops): 50 µg per mL eye drops | Primary open-angle glaucoma |

## Natamycin

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Anti-infective agents<br>◦ Local > Ophthalmological > Suspension: 5% eye drops | Infectious keratitis |

## Netilmicin

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Anti-infective agents | Therapeutic equivalent to gentamicin pour<br>Other specified conjunctivitis<br>Therapeutic equivalent to gentamicin pour<br>Infectious blepharitis |

## Ofloxacin

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Anti-infective agents<br>◦ Local > Ophthalmological > Solution (eye drops): 0.3% | Other specified conjunctivitis<br>Infectious keratitis |

## Oxytetracycline

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Anti-infective agents | Therapeutic equivalent to tetracycline pour<br>Other specified conjunctivitis<br>Therapeutic equivalent to tetracycline pour<br>Infectious keratitis<br>Therapeutic equivalent to tetracycline pour<br>Infectious blepharitis<br>Therapeutic equivalent to tetracycline pour<br>Trachoma |

## Pilocarpine

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Miotics and antiglaucoma medicines<br>◦ Local > Ophthalmological > Solution (eye drops): 2% (hydrochloride or nitrate); 4% (hydrochloride or nitrate) | Acute angle closure with pupillary block<br>Ocular hypertension<br>Primary open-angle glaucoma |

## Prednisolone

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Anti-inflammatory agents<br>◦ Local > Ophthalmological > Solution (eye drops): 0.5% (sodium phosphate) | Scleritis<br>Anterior uveitis |

## Ranibizumab

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Anti-vascular endothelial growth factor (VEGF) preparations<br>◦ . | Retinal venous occlusions<br>Diabetic macular oedema<br>Degenerative high myopia<br>Age-related macular degeneration |

## Silver nitrate

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Anti-infective agents<br>◦ Local > Ophthalmological > Solution (eye drops): 1% | Neonatal conjunctivitis or dacryocystitis |

## Sodium hyaluronate

| Section | Indications |
| --- | --- |
| Ophthalmological preparations | Therapeutic equivalent to hypromellose pour<br>Keratoconjunctivitis sicca |

## Sulfacetamide

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Anti-infective agents<br>◦ Local > Ophthalmological > Ointment: 10% (sodium salt) | Other specified conjunctivitis |

## Tetracaine

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Local anaesthetics<br>◦ Local > Ophthalmological > Solution (eye drops): 0.5% (hydrochloride) | Local anaesthetics |

## Tetracycline

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Anti-infective agents<br>◦ Local > Ophthalmological > Ointment: 1% (tetracycline hydrochloride) | Neonatal conjunctivitis or dacryocystitis<br>Other specified conjunctivitis<br>Infectious keratitis<br>Infectious blepharitis<br>Trachoma |

## Timolol

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Miotics and antiglaucoma medicines<br>◦ Local > Ophthalmological > Solution (eye drops): 0.25% (hydrogen maleate); 0.5% (hydrogen maleate) | Ocular hypertension<br>Primary open-angle glaucoma |

## Tobramycin

| Section | Indications |
| --- | --- |
| Ophthalmological preparations > Anti-infective agents | Therapeutic equivalent to gentamicin pour<br>Other specified conjunctivitis<br>Therapeutic equivalent to gentamicin pour<br>Infectious blepharitis |