Liste modèle des médicaments essentiels

Acamprosate calcium

Section	Indications
Medicines for alcohol use disorders ◦ Oral > Solid > tablet: 333 mg	Disorders due to use of alcohol

Buprenorphine

Section	Indications
Medicines for opioid use disorders	Therapeutic equivalent to methadone pour Opioid dependence

Bupropion

Section	Indications
Medicines for nicotine use disorders ◦ Oral > Solid: 150 mg sustained-release (hydrochloride)	Nicotine dependence

Cytisine (cytisinicline)

Section	Indications
Medicines for nicotine use disorders ◦ Oral > Solid > tablet: 1.5 mg	Nicotine dependence

Methadone

Section	Indications
Medicines for opioid use disorders ◦ Oral > Liquid: 5 mg per 5 mL (methadone hydrochloride); 10 mg per 5 mL (methadone hydrochloride); 5 mg per mL concentrate for oral liquid (methadone hydrochloride); 10 mg per mL concentrate for oral liquid (methadone hydrochloride)	Opioid dependence

Naltrexone

Section	Indications
Medicines for alcohol use disorders ◦ Parenteral > General injections > IM: 380 mg in vial (extended-release) ◦ Oral > Solid > tablet: 50 mg	Disorders due to use of alcohol

Nicotine replacement therapy

Section	Indications
Medicines for nicotine use disorders ◦ Local > Topical > Transdermal patch: 5 to 30 mg per 16 hour; 7 to 21 mg per 24 hour ◦ Oral > Solid > lozenge: 2 mg; 4 mg lozenge ◦ Local > Buccal > Oral spray: 1 mg per actuation ◦ Local > Buccal > Chewing gum: 2 mg; 4 mg	Nicotine dependence

Varenicline

Section	Indications
Medicines for nicotine use disorders ◦ Oral > Solid: 0.5 mg; 1 mg	Nicotine dependence