# Liste modèle des médicaments essentiels

## Aztreonam

| Section | Indications |
| --- | --- |
| Reserve group antibiotics<br>◦ Parenteral > General injections > unspecified: 1 g in vial powder for injection; 2 g in vial powder for injection | Other specified bacterial diseases |

## Cefiderocol

| Section | Indications |
| --- | --- |
| Reserve group antibiotics<br>◦ Parenteral > General injections > IV: 1 g in vial (as sulfate tosylate) powder for injection | Carbapenem-resistant Pseudomonas aeruginosa<br>Other carbapenem resistant Enterobacterales<br>Carbapenem resistant Klebsiella pneumoniae<br>Carbapenem resistant Escherichia coli |

## Ceftaroline

| Section | Indications |
| --- | --- |
| Reserve group antibiotics<br>◦ Parenteral > General injections > IV: 400 mg in vial (as fosamil) powder for injection; 600 mg in vial (as fosamil) powder for injection | Other specified bacterial diseases |

## Ceftazidime + avibactam

| Section | Indications |
| --- | --- |
| Reserve group antibiotics<br>◦ Parenteral > General injections > IV: 2 g in vial + 0.5 g in vial powder for injection | Carbapenem-resistant Pseudomonas aeruginosa<br>Other carbapenem resistant Enterobacterales<br>Carbapenem resistant Klebsiella pneumoniae<br>Carbapenem resistant Escherichia coli |

## Ceftolozane + tazobactam

| Section | Indications |
| --- | --- |
| Reserve group antibiotics<br>◦ Parenteral > General injections > IV: 1 g + 0.5 g powder for injection | Carbapenem-resistant Pseudomonas aeruginosa |

## Colistin (injection)

| Section | Indications |
| --- | --- |
| Reserve group antibiotics<br>◦ Parenteral > General injections > unspecified: 1 million IU in vial (as colistemethate sodium)<br>◦ Parenteral > General injections > unspecified: 1 million IU in vial (as colistemethate sodium) (equivalent to 34 mg colistin base activity) | Other specified bacterial diseases<br>Carbapenem resistant Acinetobacter baumannii<br>Carbapenem-resistant Pseudomonas aeruginosa<br>Other carbapenem resistant Enterobacterales<br>Carbapenem resistant Klebsiella pneumoniae<br>Carbapenem resistant Escherichia coli |

## Daptomycin

| Section | Indications |
| --- | --- |
| Reserve group antibiotics<br>◦ Parenteral > General injections > IV: 350 mg in vial powder for injection; 500 mg in vial powder for injection | Other specified bacterial diseases |

## Eravacycline

| Section | Indications |
| --- | --- |
| Reserve group antibiotics<br>◦ Parenteral > General injections > IV: 50 mg lyophilized powder for injection | Other carbapenem resistant Enterobacterales |

## Fifth generation cephalosporins

| Section | Indications |
| --- | --- |
| Reserve group antibiotics<br>◦ . | Other specified bacterial diseases |

## Fosfomycin (injection)

| Section | Indications |
| --- | --- |
| Reserve group antibiotics<br>◦ Parenteral > General injections > IV: 2 g in vial (as sodium) powder for injection; 4 g in vial (as sodium) powder for injection | Other specified bacterial diseases<br>Carbapenem resistant Acinetobacter baumannii<br>Carbapenem-resistant Pseudomonas aeruginosa<br>Other carbapenem resistant Enterobacterales |

## Imipenem + cilastatin + relebactam

| Section | Indications |
| --- | --- |
| Reserve group antibiotics<br>◦ Parenteral > General injections > IV: 500 mg + 500 mg + 250 mg powder for injection<br>◦ Parenteral > General injections > IV: 500 mg (as monohydrate) + 500 mg (as sodium) + 250 mg (as monohydrate) powder for injection | Other specified other antibiotic resistant Enterobacterales<br>Carbapenem-resistant Pseudomonas aeruginosa<br>Other carbapenem resistant Enterobacterales |

## Linezolid

| Section | Indications |
| --- | --- |
| Reserve group antibiotics<br>◦ Parenteral > General injections > IV: 2 mg per mL in 300 mL bag<br>◦ Oral > Liquid: 100 mg per 5 mL powder for oral liquid<br>◦ Oral > Solid: 400 mg tablet; 600 mg tablet<br>◦ Oral > Solid > dispersible tablet: 150 mg<br>◦ Oral > Solid > tablet: 600 mg | Other specified bacterial diseases<br>Methicillin resistant Staphylococcus aureus<br>Vancomycin resistant Staphylococcus aureus<br>Vancomycin resistant Enterococcus |

## Meropenem + vaborbactam

| Section | Indications |
| --- | --- |
| Reserve group antibiotics<br>◦ Parenteral > General injections > IV: 1 g in vial + 1 g in vial powder for injection | Carbapenem-resistant Pseudomonas aeruginosa<br>Carbapenem resistant Acinetobacter baumannii<br>Other carbapenem resistant Enterobacterales<br>Carbapenem resistant Klebsiella pneumoniae<br>Carbapenem resistant Escherichia coli |

## Omadacycline

| Section | Indications |
| --- | --- |
| Reserve group antibiotics<br>◦ Parenteral > General injections > IV: 100 mg lyophilized powder for injection<br>◦ Oral > Solid: 300 mg | Methicillin resistant Staphylococcus aureus |

## Oxazolindinones

| Section | Indications |
| --- | --- |
| Reserve group antibiotics<br>◦ . | Other specified bacterial diseases |

## Plazomicin

| Section | Indications |
| --- | --- |
| Reserve group antibiotics<br>◦ Parenteral > General injections > IV: 500 mg per 10 mL | Carbapenem resistant Acinetobacter baumannii<br>Carbapenem-resistant Pseudomonas aeruginosa<br>Other carbapenem resistant Enterobacterales<br>Carbapenem resistant Escherichia coli<br>Carbapenem resistant Klebsiella pneumoniae |

## Polymyxin B (injection)

| Section | Indications |
| --- | --- |
| Reserve group antibiotics<br>◦ Parenteral > General injections > IV: 500000 IU in vial powder for injection (equivalent to 50 mg polymyxin B base) | Carbapenem-resistant Pseudomonas aeruginosa<br>Carbapenem resistant Acinetobacter baumannii<br>Other carbapenem resistant Enterobacterales<br>Carbapenem resistant Klebsiella pneumoniae<br>Carbapenem resistant Escherichia coli |

## Polymyxins

| Section | Indications |
| --- | --- |
| Reserve group antibiotics<br>◦ . | Other specified bacterial diseases |

## Tedizolid

| Section | Indications |
| --- | --- |
| Reserve group antibiotics | Therapeutic equivalent to linezolid pour<br>Methicillin resistant Staphylococcus aureus<br>Therapeutic equivalent to linezolid pour<br>Vancomycin resistant Staphylococcus aureus<br>Therapeutic equivalent to linezolid pour<br>Vancomycin resistant Enterococcus |

## Tigecycline

| Section | Indications |
| --- | --- |
| Reserve group antibiotics<br>◦ Parenteral > General injections > IV: 50 mg in vial powder for injection | Other specified bacterial diseases |