



Liste modèle des médicaments essentiels

Beinaglutide [Informations générales](#)

Section

Hypoglycaemic agents

Indications

Therapeutic equivalent to [semaglutide](#) pour [Obesity](#)

Canagliflozin [Informations générales](#)

Section

Hypoglycaemic agents

Indications

Therapeutic equivalent to [empagliflozin](#) pour [Type 2 diabetes mellitus](#)

Dapagliflozin [Informations générales](#)

Section

Hypoglycaemic agents

Indications

Therapeutic equivalent to [empagliflozin](#) pour [Type 2 diabetes mellitus](#)

Dulaglutide [Informations générales](#)

Section

Hypoglycaemic agents

Indications

Therapeutic equivalent to [semaglutide](#) pour [Obesity](#) Therapeutic equivalent to [semaglutide](#) pour [Type 2 diabetes mellitus](#)

Empagliflozin [Informations générales](#)

Section

Hypoglycaemic agents

◦ Oral > Solid: 10 mg; 25 mg

Indications

[Type 2 diabetes mellitus](#)

Exenatide [Informations générales](#)

Section

Hypoglycaemic agents

Indications

Therapeutic equivalent to [semaglutide](#) pour [Obesity](#)

Glibenclamide [Informations générales](#)

Section

Hypoglycaemic agents

- Oral > Solid: 5 mg; 2.5 mg

Indications

[Type 2 diabetes mellitus](#)

Gliclazide [Informations générales](#)

Section

Hypoglycaemic agents

- Oral > Solid: 30 mg controlled-release; 60 mg controlled-release; 80 mg controlled-release

Indications

[Type 2 diabetes mellitus](#)

Liraglutide [Informations générales](#)

Section

Hypoglycaemic agents

Indications

Therapeutic equivalent to [semaglutide](#) pour [Obesity](#) Therapeutic equivalent to [semaglutide](#) pour [Type 2 diabetes mellitus](#)

Metformin [Informations générales](#)

Section

Hypoglycaemic agents

- Oral > Solid: 500 mg (hydrochloride)

Indications

[Type 2 diabetes mellitus](#)

Semaglutide [Informations générales](#)

Section

Hypoglycaemic agents

- Parenteral > General injections > SC: 0.25 mg in pre-filled pen injection solution; 0.5 mg in pre-filled pen injection solution; 1 mg in pre-filled pen injection solution; 2 mg in pre-filled pen injection solution; 4 mg in pre-filled pen injection solution
- Oral > Solid > tablet: 3 mg
- Parenteral > General injections > SC: 0.68 mg per mL; 1.34 mg per mL; 2.68 mg per mL

Indications

[Obesity](#) [Type 2 diabetes mellitus](#)

Tirzepatide [Informations générales](#)

Section

Hypoglycaemic agents

Indications

Therapeutic equivalent to [semaglutide](#) pour [Obesity](#) Therapeutic equivalent to [semaglutide](#) pour [Type 2 diabetes mellitus](#)

Tolbutamide [Informations générales](#)

Section

Hypoglycaemic agents

- Oral > Solid: 500 mg

Indications

[Type 2 diabetes mellitus](#)