

## [Glibenclamide](#)

Not recommended as an essential medicine

Section:

[18. Medicines for endocrine disorders 18.5. Medicines for diabetes 18.5.2. Hypoglycaemic agents](#)

Codes ATC: [A10BB01](#)

Indication

Type 2 diabetes mellitus Code ICD11: [5A11](#)

INN

Glibenclamide

Type de médicament

Chemical agent

Type de liste

Liste de base

Formulations

**Oral > Solid:** 5 mg ; 2.5 mg

Historique des statuts LME

Ajouté pour la première fois en 1982 ([TRS 685](#))

Modifié en 1984 ([TRS 722](#))

Retiré en 1987 ([TRS 770](#))

Ajouté en 1997 ([TRS 882](#))

Retiré en 2013 ([TRS 985](#))

Sexe

Tous

Âge

Adolescents et adultes

Équivalence thérapeutique

La recommandation concerne ce médicament spécifique

Renseignements sur le brevet

Patents have expired in most jurisdictions

Lire la suite [sur les brevets.](#)

Wikipédia

[Glibenclamide](#)

DrugBank

[Glibenclamide \(Glyburide\)](#)

Résumé des preuves et recommandation du comité d'experts



At its 18th meeting, the Expert Committee requested a review on the safety of sulfonylureas in elderly patients to determine whether updates to the EML were needed. The review was prepared by Mr Harinder Singh Chahal, Doctor of Pharmacy Candidate at the University of California - San Francisco, San Francisco, CA, USA. Expert reviews were prepared by Mr Andrew Gray and Dr Kuruvilla Prasad Mathews. Comments were received from Dr Shanthi Mendis, Director, Management of Noncommunicable Diseases, WHO, and Dr Myriam Henkens, International Medical Coordinator, Médecins Sans Frontières. The estimated worldwide prevalence of diabetes in the elderly population (60 years and above) is 18.6% or 134.6 million people (1). In 2013 the Expert Committee noted that the EML listed only glibenclamide from the sulfonylurea category. Elderly patients are a significant proportion of type II diabetics and the safety of sulfonylureas in this group is therefore important. The review evaluated the comparative safety and efficacy of four second-generation sulfonylureas for the treatment of type 2 non-insulin dependent diabetes in elderly patients. The medications reviewed were glibenclamide (also called glyburide), gliclazide, glimepiride and glipizide. The review also analysed the cost of the four medications as well as their availability on the national essential medicines lists of 40 low- and middle-income countries. A 2007 meta-analysis of 21 studies showed that, on the basis of HbA1c results, glibenclamide compared with other sulfonylureas - including gliclazide, glimepiride and glipizide - did not have an increased efficacy in the treatment of diabetes (2). The same meta-analysis showed that there was an increased risk of hypoglycaemia of 52% with glibenclamide when compared with other insulin-secreting anti-diabetes medicines and an 83% higher risk compared with other sulfonylureas (2). A retrospective cohort study of more than 13 000 patients concluded that glibenclamide had the highest rate of hypoglycaemia at 16.9 per 1000 person-years, compared with all other sulfonylureas (3). The authors also concluded that the physiological changes associated with increasing age, such as declining renal and hepatic function, as well as polypharmacy and concurrent illnesses, additionally predispose elderly patients to hypoglycaemia; this predisposition is further compounded by the use of glibenclamide. Another retrospective, cohort study of more than 33 000 patients in the United Kingdom showed that the risk of hypoglycaemia was higher with glibenclamide when compared with other sulfonylureas (4). The authors also concluded that patients older than 65 years were at higher risk of hypoglycaemia compared with adults less than 65 years of age with a relative risk of 1.27 (95% CI: 1.06-1.51). In the analysis of the national essential medicines lists, the most widely available second-generation sulfonylurea was glibenclamide, with an overall listing on 39 of the 40 national essential medicines lists (97.5%), followed by gliclazide and glipizide which were available on 50% and 27.5% of the national lists, respectively. The Expert Committee discussed the fact that all four sulfonylureas are available as generics, but there is considerable variation in price between countries and therefore it is not possible to make a clear decision on the basis of cost. However, glibenclamide appeared to be the cheapest in most countries. The Expert Committee decided that glibenclamide should be replaced in the core list with gliclazide (30 mg, 60 mg, 80 mg), with a square box symbol as the example of a second-generation sulfonylurea, and that a note should be added to the effect that glibenclamide should not be used in patients aged 60 years and older. References: 1. IDF diabetes atlas, 6th edition. Brussels: International Diabetes Federation; 2013 (<http://www.idf.org/diabetesatlas/6e/diabetes>, accessed 27 November 2013). 2. Gangji AS, Cukierman T, Gerstein HC, Goldsmith CH, Clase CM. A systematic review and meta-analysis of hypoglycemia and cardiovascular events: a comparison of glyburide with other secretagogues and with insulin. *Diabetes Care*. 2007;30(2):389-94. <http://dx.doi.org/10.2337/dc06-1789> PMID:17259518 3. Shorr RI, Ray WA, Daugherty JR, Gri n MR. Individual sulfonylureas and serious hypoglycemia in older people. *J Am Geriatr Soc*. 1996;44(7):751-5. PMID:8675920

4. van Staa T, Abenhaim L, Monette J. Rates of hypoglycemia in users of sulfonylureas. *J Clin Epidemiol.* 1997;50(6):735-41. [http://dx.doi.org/10.1016/S0895-4356\(97\)00024-3](http://dx.doi.org/10.1016/S0895-4356(97)00024-3) PMID:9250272