





EMLc
Codes ATC: A04AD12

Indication	Nausea or vomiting <span>Code ICD11: MG10</span>
INN	Aprepitant
Type de médicament	Chemical agent
Type de liste	Liste complémentaire (EML) (EMLc)
Formulations	Oral > Liquid: 125 mg powder for oral suspension (in sachet) Oral > Solid > capsule: 80 mg (EML) ; 125 mg (EML) ; 165 mg (EML)
Historique des statuts LME	Ajouté pour la première fois en 2019 (TRS 1021) Modifié en 2025 (TRS 1064)
Sexe	Tous
Âge	Aussi recommandé pour les enfants
Équivalence thérapeutique	La recommandation concerne ce médicament spécifique
Renseignements sur le brevet	Main patents have expired but secondary patents might remain active in some jurisdictions. For more information on specific patents and license status for developing countries visit <a href="http://www.MedsPal.org">www.MedsPal.org</a>  Lire la suite <a href="#">sur les brevets.</a> 
Wikipédia	<a href="#">Aprepitant</a> 
DrugBank	<a href="#">Aprepitant</a> 

## Résumé des preuves et recommandation du comité d'experts

Following the review of the age-appropriateness of formulations of essential medicines for children, the Expert Committee recommended the removal from the EMLc of aprepitant capsules (80 mg, 125 mg and 165 mg). These formulations cannot deliver the recommended dose of aprepitant for children aged below 12 years.

