Oral rehydration salts

EMLc | Codes ATC: A07CA

Indication	Diarrhoea Code ICD11: ME05.1
Type de médicament	Chemical agent
Type de liste	Liste de base
Additional notes	Trisodium citrate dihydrate may be replaced by sodium hydrogen carbonate (sodium bicarbonate) 2.5 g/L. However, as the stability of the latter formulation is very poor under tropical conditions, it is recommended only when manufactured for immediate use.
Formulations	Powder for dilution in 200 mL, 500 mL, 1 L: - glucose 75 mEq or mmol/L - sodium 75 mEq or mmol/L - chloride 65 mEq or mmol/L - potassium 20 mEq or mmol/L - citrate 10 mEq or mmol/L - osmolarity 245 mOsm/L - glucose 13.5 g/L - sodium chloride 2.6 g/L - potassium chloride 1.5 g/L - trisodium citrate dihydrate 2.9 g/L
Historique des statuts LME	Ajouté pour la première fois en 1977 (TRS 615) Modifié en 1984 (TRS 722) Modifié en 2003 (TRS 920) Modifié en 2007 (TRS 950) Modifié en 2025 (TRS 1064)
Sexe	Tous
Âge	Aussi recommandé pour les enfants
Équivalence thérapeutique	La recommandation concerne ce médicament spécifique
Renseignements sur le brevet	Patents have expired in most jurisdictions Lire la suite sur les brevets.
Wikipédia	Oral rehydration salts

Résumé des preuves et recommandation du comité d'experts

Following the review of the age-appropriateness of formulations for children, the Expert Committee recommended:

- to modify the formulation description to better describe the available dosage forms.