




Section: 26. Solutions correcting water, electrolyte and acid–base disturbances > 26.2. Solutions correcting water, electrolyte and acid-base disturbances > Parenteral

		<b>EMLc</b>	ATC codes: <b>B05CX01</b>
<b>Indication</b>	Other specified disorders of fluid, electrolyte or acid-base balance	ICD11 code: <b>5C5Y</b>	
<b>Medicine type</b>	Chemical agent		
<b>List type</b>	Core		
<b>Formulations</b>	Parenteral > General injections > IV: 5% isotonic ; 10% hypertonic		
<b>EML status history</b>	First added in 1977 ( <a href="#">TRS 615</a> ) Changed in 1979 ( <a href="#">TRS 641</a> ) Changed in 1999 ( <a href="#">TRS 895</a> ) Changed in 2007 ( <a href="#">TRS 950</a> ) Changed in 2009 ( <a href="#">TRS 958</a> )		
<b>Sex</b>	All		
<b>Age</b>	Also recommended for children		
<b>Therapeutic alternatives</b>	The recommendation is for this specific medicine		
<b>Patent information</b>	Patents have expired in most jurisdictions Read more <a href="#">about patents</a> . 		
<b>Wikipedia</b>	<a href="#">Glucose</a> 		
<b>DrugBank</b>	<a href="#">Glucose</a> 		

### Summary of evidence and Expert Committee recommendations

Formulation correction to specify 10% glucose solution as hypertonic.

