

Section: 26. Solutions correcting water, electrolyte and acid-base disturbances > 26.2. Solutions correcting water, electrolyte and acid-base disturbances > Parenteral

| | | EMLc | ATC codes: Pending |
|--------------------------|---|------|--------------------|
| Indication | Other specified disorders of fluid, electrolyte or acid-base balance | | ICD11 code: 5C5Y |
| Medicine type | Chemical agent | | |
| List type | Core (EML) (EMLc) | | |
| Formulations | Parenteral > General injections > IV: 4% + 0.18% (equivalent to Na+ 30 mmol/L and Cl- 30 mmol/L) (EML); 5% + 0.9% (equivalent to Na+ 150 mmol/L and Cl- 150 mmol/L) (EMLc); 5% + 0.45% (equivalent to Na+ 75 mmol/L and Cl- 75 mmol/L) (EMLc) | | |
| EML status history | First added in 1979 (TRS 641) Changed in 2007 (TRS 950) Changed in 2009 (TRS 958) | | |
| Sex | All | | |
| Age | Also recommended for children | | |
| Therapeutic alternatives | The recommendation is for this specific medicine | | |
| Patent information | Patents have expired in most jurisdictions Read more about patents.  | | |
| Wikipedia | Glucose + sodium chloride  | | |
| DrugBank | Glucose  , Sodium chloride  | | |

Summary of evidence and Expert Committee recommendations

The EMLc Subcommittee recommended the deletion of glucose + sodium chloride 4% + 0.18% injectable solution from the EMLc. It remains listed on the EML.

