





Codes ATC: J01MA01

Code ICD11: ML32.00

| | |
|------------------------------|--|
| Indication | Multi-drug resistant Mycobacterium tuberculosis |
| INN | Ofloxacin |
| Type de médicament | Chemical agent |
| Groupes d'antibiotiques |  WATCH |
| Type de liste | Liste complémentaire |
| Additional notes | Levofloxacin may be an alternative based on availability and programme considerations. |
| Formulations | Oral > Solid: 200 mg ; 400 mg |
| Historique des statuts LME | Ajouté pour la première fois en 1999 (TRS 895) Modifié en 2007 (TRS 946) Retiré en 2013 (TRS 985) |
| Sexe | Tous |
| Âge | Adolescents et adultes |
| Équivalence thérapeutique | La recommandation concerne ce médicament spécifique |
| Renseignements sur le brevet | Patents have expired in most jurisdictions Lire la suite sur les brevets.  |
| Wikipédia | Ofloxacin  |
| DrugBank | Ofloxacin  |

Résumé des preuves et recommandation du comité d'experts

On the basis of WHO guideline recommendations, public health needs and available evidence on efficacy and safety, the Expert Committee recommended that ofloxacin should be replaced by levofloxacin, with a note to indicate that ofloxacin and moxifloxacin may be used as alternatives in the treatment of MDR-TB.

