Erythromycin

NOT RECOMMENDED AS AN ESSENTIAL MEDICINE

Section: 6. Anti-infective medicines > 6.2. Antibacterials > 6.2.2. Watch group antibiotics

EMLc | Codes ATC: J01FA01

Indication	Bacterial infection of unspecified site Code ICD11: 1C41
INN	Erythromycin
Type de médicament	Chemical agent
Groupes d'antibiotiques	WATCH
Type de liste	Liste de base
Formulations	**Parenteral > General injections > IV:** 500 mg in vial powder for injection (as lactobionate) **Oral > Liquid:** 125 mg per 5 mL (as stearate or ethylsuccinate) **Oral > Solid:** 250 mg (as stearate or ethylsuccinate)
Historique des statuts LME	Ajouté pour la première fois en 1977 (TRS 615) Modifié en 1979 (TRS 641) Modifié en 1987 (TRS 770) Modifié en 2007 (TRS 950) Retiré en 2017 (TRS 1006)
Sexe	Tous
Âge	Aussi recommandé pour les enfants
Équivalence thérapeutique	Des médicaments appartenant à la même classe pharmacologique peuvent être utilisés
Limites de l'équivalence thérapeutique	The square box only applies to the listing of erythromycin on the EML
Limites d'équivalence thérapeutique pour l'EMLc	The square box does not apply to the listing of erythromycin on the EMLc
Renseignements sur le brevet	Patents have expired in most jurisdictions Lire la suite sur les brevets.
Wikipédia	Erythromycin
DrugBank	Erythromycin

Résumé des preuves et recommandation du comité d'experts

Erythromycin was removed from the EML and EMLc following the comprehensive review of antibiotics by the Expert Committee in 2017. However, it is noted in the Model Lists as an alternative to clarithromycin for treatment of pharyngitis