

| | | | |
|------------------------------|---|----------------------------|---------------------------|
| | | EMLc | Codes ATC: J01DH51 |
| Indication | Pseudomonas aeruginosa resistant to other antibiotic | Code ICD11: MG50.8Y | |
| INN | Imipenem + cilastatin | | |
| Type de médicament | Chemical agent | | |
| Groupes d'antibiotiques | W WATCH | | |
| Type de liste | Liste complémentaire (EML) (EMLc) | | |
| Formulations | Parenteral > General injections > IV: 250 mg (as monohydrate) + 250 mg (as sodium salt) powder for injection ; 500 mg (as monohydrate) + 500 mg (as sodium salt) powder for injection | | |
| Historique des statuts LME | Ajouté pour la première fois en 1997 (TRS 882) Modifié en 2002 (TRS 914) Modifié en 2007 (TRS 950) Modifié en 2009 (TRS 958) Retiré en 2017 (TRS 1006) | | |
| Sexe | Tous | | |
| Âge | Aussi recommandé pour les enfants | | |
| Équivalence thérapeutique | La recommandation concerne ce médicament spécifique | | |
| Renseignements sur le brevet | Patents have expired in most jurisdictions Lire la suite sur les brevets. ↗ | | |
| Wikipédia | Imipenem + cilastatin ↗ | | |
| DrugBank | Imipenem ↗ Cilastatin ↗ | | |

Résumé des preuves et recommandation du comité d'experts

As part of the syndrome-based comprehensive review of antibacterials on the Model Lists, imipenem + cilastatin was removed from the EML and EMLc. However, it is acknowledged in the lists as an alternative to meropenem except for acute bacterial meningitis where meropenem is preferred.

