Ethionamide

Section: 6. Anti-infective medicines > 6.2. Antibacterials > 6.2.5. Antituberculosis medicines

EMLc Codes ATC: J04AD03

Indication	Tuberculous meningitis Code ICD11: 1B11.0
INN	Ethionamide
Type de médicament	Chemical agent
Type de liste	Liste de base
Formulations	**Oral > Solid > dispersible tablet:** 125 mg (EMLc) **Oral > Solid > tablet:** 250 mg
Historique des statuts LME	Ajouté pour la première fois en 2023 (TRS 1049)
Sexe	Tous
Âge	Aussi recommandé pour les enfants
Équivalence thérapeutique	La recommandation concerne ce médicament spécifique
Renseignements sur le brevet	Patents have expired in most jurisdictions Lire la suite sur les brevets.
Wikipédia	Ethionamide
DrugBank	Ethionamide

Recommandation du comité d'experts

The Expert Committee noted that tuberculosis meningitis is responsible for considerable morbidity and mortality, and that a shorter, intensified ethionamide-containing treatment regimen in children and adolescents has shown favourable outcomes in comparison with the alternative WHO-recommended 12-month regimen of isoniazid, rifampicin, pyrazinamide and ethambutol. The Committee therefore recommended the inclusion of ethionamide on the core list of the EML and EMLc for the new indication of drug-susceptible tuberculosis meningitis in children and adolescents, consistent with the recommendations in current WHO guidelines for management of drug-susceptible tuberculosis meningitis in children and adolescents.

Contexte

Ethionamide is currently included on the complementary list of the EML and EMLc for use in the treatment of multidrug-resistant tuberculosis.

Pertinence pour la santé publique

Tuberculous meningitis is the most lethal form of tuberculosis. Globally in 2019, there were an estimated 164 000 cases and 78 200 deaths due to tuberculous meningitis (1). Mortality and severe permanent disabilities remain high in both children and adults, particularly in people living with HIV (2,3).

Bénéfices

The application referenced a systematic review and meta-analysis of seven cohort studies comparing the

effectiveness of shorter regimens including at a minimum isoniazid, rifampicin and pyrazinamide, versus the WHO-recommended 12-month regimen of isoniazid, rifampicin, ethambutol and pyrazinamide in children and adolescents with drug-susceptible tuberculosis meningitis (4). This meta-analysis informed a 2022 WHO guideline recommendation in favour of the shorter regimen (conditional recommendation; very low-certainty evidence). Details of the findings of the systematic review were not provided in the application but are summarized below. Three of the included studies (724 patients) evaluated a 6-month intensive regimen of isoniazid, rifampicin, pyrazinamide and ethionamide. This regimen was associated with a lower pooled proportion of death (5.5%, 95% confidence interval (CI) 2.1% to 13.4%) compared with the 12-month regimen (23.9%, 95% CI 17.5% to 31.7%). The pooled proportions of treatment success were 94.6% (95% CI 73.9% to 99.1%) for the 6-month intensive regimen and 75.4% (95% CI 68.7% to 81.1%) for the 12-month regimen. For survivors who completed treatment and who had neurological sequalae the pooled proportions were 66.0% (95% CI 55.3% to 75.3%) for the 6-month regimen and 36.3% (95% CI 30.1% to 43.0%) for the 12-month regimen, although there was substantial heterogeneity for both regimens. For survivors who completed treatment and who did not have neurological sequalae, the pooled proportions were 29.9% (95% CI 20.4% to 41.4%) and 47.9% (95% CI 42.1% to 53.7%) for the 6-month and 12-month regimens, respectively.

Torts

Harms associated with the use of ethionamide were not discussed in the application. From one of the studies included in the systematic review that evaluated 6- and 9-month intensified regimens of isoniazid, rifampicin, pyrazinamide and ethionamide in children with tuberculosis meningitis, treatment-induced hepatotoxicity was reported in 5% of the children (5).

Rapport coût/efficacité

No information was provided in the application. The 2023 Global Drug Facility catalogue reports the price of ethionamide 250 mg tablets as US$ 9.16 for 100 tablets, and of ethionamide 125 mg dispersible tablets as US$ 13.30–14.48 for 100 tablets.

Directives de l'OMS

Current WHO guidelines for the management of tuberculosis in children and adolescents include a conditional recommendation (very low-certainty evidence) that in children and adolescents with bacteriologically confirmed or clinically diagnosed tuberculosis meningitis (without suspicion or evidence of multidrug- or rifampicin-resistant tuberculosis), a 6-month intensive regimen of isoniazid, rifampicin, pyrazinamide and ethionamide may be used as an alternative to the 12-month regimen of isoniazid, rifampicin, ethambutol and pyrazinamide (6).

Disponibilité

Ethionamide tablets and dispersible tablets are available through the Stop TB Partnership’s Global Drug Facility.

Autres considérations

In adults, WHO guidelines recommend that drug-susceptible tuberculosis meningitis be treated with the same regimen used for pulmonary tuberculosis, that is, a 6-month regimen composed of 2 months of isoniazid, rifampicin, pyrazinamide and ethambutol, followed by 4 months of isoniazid and rifampicin, noting that some expert groups suggest longer therapy (7).

1. Dodd PJ OM, Cresswell FV, Stadelman AM, Lan NH, Thuong NTT, et al. The global burden of tuberculous meningiti s in adults: a modelling study. PLOS Glob Public Health. 2021;1(12):e0000069.
2. Stadelman AM, Ellis J, Samuels THA, Mutengesa E, Dobbin J, Ssebambulidde K, et al. Treatment outcomes in adult tuberculous meningitis: a systematic review and meta-analysis. Open Forum Infect Dis. 2020;7(8):ofaa257.
3. Chiang SS, Khan FA, Milstein MB, Tolman AW, Benedetti A, Starke JR, et al. Treatment outcomes of childhood tub erculous meningitis: a systematic review and meta-analysis. Lancet Infect Dis. 2014;14(10):947–57.
4. Sulis G, Tavaziva G, Gore G, Benedetti A, Solomons R, van Toorn R, et al. Comparative effectiveness of regimens for drug-susceptible tuberculous meningitis in children and adolescents: a systematic review and aggregate-level d

ata meta-analysis. Open Forum Infect Dis. 2022;9(6):ofac108.
5. van Toorn R, Schaaf HS, Laubscher JA, van Elsland SL, Donald PR, Schoeman JF. Short intensified treatment in children with drug-susceptible tuberculous meningitis. Pediatr Infect Dis J. 2014;33(3):248–52.
6. WHO consolidated guidelines on tuberculosis: module 5: management of tuberculosis in children and adolescents. Geneva: World Health Organization; 2022 (https://apps.who.int/iris/handle/10665/352522, accessed 6 October 2023).
7. WHO consolidated guidelines on tuberculosis: module 4: treatment: drug-susceptible tuberculosis treatment. Geneva: World Health Organization; 2022 (https://apps.who.int/iris/handle/10665/353829, accessed 6 October 2023).

