

Codes ATC: **N02BA01**

Indication	Pain <span style="background-color: #0070C0; color: white; padding: 2px;">Code ICD11: <b>ML0Z</b></span>
Type de médicament	Chemical agent
Type de liste	Liste de base
Formulations	Oral > Solid: 100 to 500 mg Local > Rectal > Suppository: 50 to 150 mg
Historique des statuts LME	Ajouté pour la première fois en 1977 ( <a href="#">TRS 615</a> )
Sexe	Tous
Âge	Adolescents et adultes
Équivalence thérapeutique	La recommandation concerne ce médicament spécifique
Renseignements sur le brevet	Patents have expired in most jurisdictions Lire la suite <a href="#">sur les brevets.</a>
Wikipédia	<a href="#">Acetylsalicylic acid</a>
DrugBank	<a href="#">Acetylsalicylic acid</a>

### Résumé des preuves et recommandation du comité d'experts

Acetylsalicylic acid was included on the first EML in 1977 as an analgesic/antipyretic/non-steroidal anti-inflammatory medicine. It was included in the complementary list of the first EMLc in 2007 specifically for use in rheumatic fever, juvenile arthritis, and Kawasaki disease. The EMLc Subcommittee agreed that the management of pain in children is an essential clinical need but that aspirin is not recommended for use in children under 16 years in many settings, and had only a limited role, if any, in the management of inflammatory conditions in children. This listing was deleted in 2013, and a new section (Section 30: Medicines for diseases of joints) was created which included acetylsalicylic acid for juvenile joint diseases.

