

## [Hypromellose](#)

Le Comité d'experts, après évaluation, refuse d'inscrire le médicament proposé dans la demande.

La Liste Modèle des Médicaments Essentiels fait état des raisons que les membres du Comité ont identifiées pour refuser l'inscription.

Refusée

Section:

[21. Ophthalmological preparations](#)

Codes ATC: [S01KA02](#)

EMLc

Indication

Keratoconjunctivitis sicca Code ICD11: [9A79](#)

INN

Hypromellose

Type de médicament

Diagnostic agent

Type de liste

Liste de base

Formulations

**Local > Ophthalmological > Solution (eye drops): 0.3%**

Historique des statuts LME

Demande refusée en 2023 ([TRS 1049](#))

Sexe

Tous

Âge

Aussi recommandé pour les enfants

Équivalence thérapeutique

[carmellose](#)

[sodium hyaluronate](#) (Codes ATC: [S01KA01](#))

Renseignements sur le brevet

Patents have expired in most jurisdictions

Lire la suite [sur les brevets](#).

Wikipédia

[Hypromellose](#)

DrugBank

[Hypromellose](#)

Recommandation du comité d'experts



The Committee noted that dry eye disease was a chronic and progressive condition and a common reason for ophthalmic outpatient visits. Severe dry eye disease, if untreated, can lead to ocular infection and inflammation, corneal abrasions, corneal ulcers and vision loss. Based on the evidence presented, the Committee accepted that hypromellose was a generally safe and effective ocular surface lubricant for reducing the signs and symptoms of dry eye disease for patients with mild to moderate disease. Its effectiveness and safety are comparable to other artificial tears preparations. However, the Committee noted that the available data were limited by the variable definition of dry eye disease applied in published studies and the disease severity examined, and that compliance with treatment was rarely quantified. As a result, the optimal composition, dose, formulation or formulations for artificial tears preparations for the treatment of dry eye disease have not been demonstrated. The Committee also considered that the sight-threatening complications of dry eye disease were primarily associated with severe forms of the condition. Limited evidence was available of the effectiveness of hypromellose for improving relevant clinical outcomes compared with other artificial tear preparations, including combinations, specifically in patients with severe dry eye disease. The Expert Committee therefore did not recommend inclusion of hypromellose on the EML and EMLc for the treatment of dry eye disease in adults and children.

Contexte



Artificial tears preparations for the treatment of dry eye disease have not previously been considered for inclusion on the Model Lists.

Pertinence pour la santé publique



Dry eye disease is a multifactorial disease of the ocular surface that is characterized by a loss of homeostasis of the tear film. It is accompanied by ocular symptoms, such as visual disturbance and discomfort (1,2). The negative impact on vision can limit education and work productivity by interfering with patients' daily activities, such as sustained visual attention when reading, writing, driving and using digital display monitors (3,4). Patients with dry eye disease also report psychological concerns and higher levels of anxiety and depression compared with those without dry eyes (5). Risk factors for dry eye disease include age 50 years or older (6,7), female sex (8,9), wearing contact lenses or a history of refractive surgery (10), exposure to environments with low relative humidity and extremes of temperature (10), certain chronic and autoimmune conditions (10,11), medication use (9,12) and prolonged engagement in visual tasks (13,14). Dry eye disease has a global prevalence ranging from about 5% to 50%, corresponding to 385 million to 3.85 billion people worldwide (15). The highest prevalence of dry eye disease has been reported in the WHO's African region, followed by the Eastern Mediterranean Region, South-East Asia Region, Western Pacific Region, European Region, and Region of the Americas (16).

Bénéfices



A Cochrane systematic review published in 2016 of 43 randomized controlled trials (3497 participants) evaluated the effectiveness of over-the-counter artificial tear applications in treating dry eye disease compared with no treatment, placebo or another class of over-the-counter artificial tears (17). The authors considered participant symptoms

(subjective) to be the primary outcome for the review. Secondary outcomes included objective measures of effectiveness (e.g. tests of vision or tear stability). Overall, the review found uncertainty with regard to whether different over-the-counter artificial tears provided similar relief of dry eye disease compared with each other or placebo, with most of the included studies producing contradictory between-group results, or no between-group differences. The quality of the evidence was judged as low due to high risks of bias and poor reporting of outcome measures. The authors concluded that over-the-counter artificial tears may be a safe and effective treatment for dry eye disease, with the literature indicating that most products have similar efficacy. A systematic review published in 2009 of 33 studies (1293 participants) assessed the efficacy of dry eye treatments with artificial tears or ocular lubricants using scoring of rose bengal stains as the outcome measure (18). Mean baseline and 30-day post-treatment scores were calculated, along with the net change and the percentage change in the rose bengal scores. A statistically significant reduction in mean rose bengal scores was observed from baseline to 30-days post-treatment with any type of artificial tears or ocular lubricant from 4.2 (standard deviation (SD) 1.6) to 2.8 (SD 1.2). The net reductions in mean rose bengal scores were -1.1 (SD 0.8) for traditional artificial tears (e.g. hypromellose), -1.2 (SD 0.7) for carbomer gels and -2.1 (SD 0.9) for hyaluronic acid-based products. No significant difference was found between traditional artificial tears and carbomer gels, but there was a significant difference between traditional artificial tears and hyaluronic acid-based products. A multiple analysis of variance (ANOVA) test, comparing outcomes using the different treatments, found no significant difference between the three groups. Across all studies, the overall net reduction in rose bengal staining after 4 weeks of treatment was 33%. The authors noted heavily skewed data for some treatments, so determined a 25% improvement in rose bengal staining scores with 1 month of treatment was more reasonable. No information was provided in the application on what constituted a clinically meaningful improvement. The application also presented brief summaries of findings of individual clinical trials comparing hypromellose artificial tears with other artificial tears, placebo or no treatment (19–27). The outcome assessed to evaluate the effectiveness of hypromellose tears was the relief of dry eye symptoms. Both hypromellose and comparator artificial tears products were generally found to be effective in relieving symptoms of dry eye disease. Most of these studies were included in the above-mentioned Cochrane systematic review (17).

#### Torts



The application stated that overall the clinical evidence surveyed suggested that hypromellose was generally safe, with occasional transient burning and stinging of the eyes. Similar levels of adverse effects were observed when hypromellose was compared with other types of artificial tears and dry-eye treatments. The Cochrane systematic review found that the use of artificial tears was relatively safe, although not without adverse events. The most common adverse events were blurred vision, ocular discomfort and foreign body sensation (17).

#### Rapport coût/efficacité



The application stated that hypromellose has been found to be cost-effective in several national studies (not referenced in the application) as it is a relatively cheap and effective treatment with considerable potential to reduce the burden on society from dry eye disease. In the United Kingdom, the price for a 10 mL bottle of preserved hypromellose 0.3% artificial tears (about 200 drops of 0.05 mL) was reported as US\$ 1.79, equivalent to an annual treatment cost of US\$ 18.37 assuming average usage of 5.7 drops per day. Similar costs were reported in Singapore and the United States, with a price per bottle of US\$ 1.52. Non-preserved and single unit-dose preparations of artificial tears are more costly to manufacture and to purchase. They may be less convenient to use than preserved and bottled preparations (29). In Singapore, the mean unit cost of preservative-containing lubricants was around US\$ 5.50, meanwhile that for preservative-free lubricants was US\$ 12.96 (30).

#### Directives de l'OMS



WHO guidelines for treatment of dry eye disease are not currently available. The 2019 WHO World report on vision recognizes that eye conditions that do not typically cause vision impairment, such as dry eye disease and conjunctivitis, are frequently among the leading reasons for patients to present to eye care services globally, and should not be overlooked (28).

#### Disponibilité



Artificial tears preparations, including hypromellose, are available on the market globally. They are produced by multiple manufacturers and are often available over the counter.

#### Afficher les références Masquer les références

1. Craig JP, Nichols KK, Akpek EK, Caffery B, Dua HS, Joo CK, et al. TFOS DEWS II definition and classification report. *Ocul Surf.* 2017;15(3):276–83.
2. The definition and classification of dry eye disease: report of the Definition and Classification Subcommittee of the International Dry Eye WorkShop (2007). *Ocul Surf.* 2007;5(2):75–92.
3. Miljanović B, Dana R, Sullivan DA, Schaumberg DA. Impact of dry eye syndrome on vision-related quality of life. *Am J Ophthalmol.* 2007;143(3):409–15.
4. Mertzanis P, Abetz L, Rajagopalan K, Espindle D, Chalmers R, Snyder C, et al. The relative burden of dry eye in patients' lives: comparisons to a U.S. normative sample. *Invest Ophthalmol Vis Sci.* 2005;46(1):46–50.
5. Li M, Gong L, Sun X, Chapin WJ. Anxiety and depression in patients with dry eye syndrome. *Curr Eye Res.* 2011;36(1):1–7.
6. Moss SE, Klein R, Klein BE. Prevalence of and risk factors for dry eye syndrome. *Arch Ophthalmol.* 2000;118(9):1264–8.
7. Rusciano D, Pezzino S, Olivieri M, Cristaldi M, Gagliano C, Lupo G, et al. Age-related dry eye lactoferrin and lactobionic acid. *Ophthalmic Res.* 2018;60(2):94–9.
8. Schaumberg DA, Sullivan DA, Buring JE, Dana MR. Prevalence of dry eye syndrome among US women. *Am J Ophthalmol.* 2003;136(2):318–26.
9. Sullivan DA, Rocha EM, Aragona P, Clayton JA, Ding J, Golebiowski B, et al. TFOS DEWS II sex, gender, and hormones report. *Ocul Surf.* 2017;15(3):284–333.
10. Milner MS, Beckman KA, Luchs JI, Allen QB, Awdeh RM, Berdahl J, et al. Dysfunctional tear syndrome: dry eye disease and associated tear film disorders – new strategies for diagnosis and treatment. *Curr Opin Ophthalmol.* 2017;27(Suppl 1):3–47.
11. Rouen PA, White ML. Dry eye disease: prevalence, assessment, and management. *Home Healthc Now.* 2018;36(2):74–83.
12. Moss SE, Klein R, Klein BE. Long-term incidence of dry eye in an older population. *Optom Vis Sci.* 2008;85(8):668–74.
13. Vehof J, Kozareva D, Hysi PG, Hammond CJ. Prevalence and risk factors of dry eye disease in a British female cohort. *Br J Ophthalmol.* 2014;98(12):1712–7.
14. Talens-Estarellles C, Sanchis-Jurado V, Esteve-Taboada JJ, Pons Á M, García-Lázaro S. How do different digital displays affect the ocular surface? *Optom Vis Sci.* 2020;97(12):1070–9.
15. Stapleton F, Alves M, Bunya VY, Jalbert I, Lekhanont K, Malet F, et al.

TFOS DEWS II epidemiology report. *Ocul Surf.* 2017;15(3):334-65. 16. Wan Y, Zhang M, Li X. The global prevalence of dry eye disease and its association with economy: a systematic review [preprint]. Durham, NC: Research Square; 2019. 17. Pucker AD, Ng SM, Nichols JJ. Over the counter (OTC) artificial tear drops for dry eye syndrome. *Cochrane Database Syst Rev.* 2016;2(2):CD009729. 18. Doughty MJ, Glavin S. Efficacy of different dry eye treatments with artificial tears or ocular lubricants: a systematic review. *Ophthalmic Physiol Opt.* 2009;29(6):573-83. 19. Boisjoly H, Charest M. Comparison of GenTeal Gel and Refresh Liquigel for persistent dry eye. *Clin Surg Ophthalmol.* 2003;21(10):430-2. 20. Donshik PC, Nelson JD, Abelson M, McCulley JP, Beasley C, Laibovitz RA. Effectiveness of BION tears, Cellufresh, Aquasite, and Refresh Plus for moderate to severe dry eye. *Adv Exp Med Biol.* 1998;438:753-60. 21. Garcia-Lázaro S, Belda-Salmerón L, Ferrer-Blasco T, Cerviño A, Montés-Micó R. Comparison of two artificial tear formulations for dry eye through high-resolution optical coherence tomography. *Clin Exp Optom.* 2011;94(6):549-56. 22. Grene RB, Lankston P, Mordaunt J, Harrold M, Gwon A, Jones R. Unpreserved carboxymethylcellulose artificial tears evaluated in patients with keratoconjunctivitis sicca. *Cornea.* 1992;11(4):294-301. 23. Iester M, Orsoni GJ, Gamba G, Taffara M, Mangiafico P, Giuffrida S, et al. Improvement of the ocular surface using hypotonic 0.4% hyaluronic acid drops in keratoconjunctivitis sicca. *Eye (Lond).* 2000;14(Pt 6):892-8. 24. Khanal S, Tomlinson A, Pearce EI, Simmons PA. Effect of an oil-in-water emulsion on the tear physiology of patients with mild to moderate dry eye. *Cornea.* 2007;26(2):175-81. 25. Wang IJ, Lin IC, Hou YC, Hu FR. A comparison of the effect of carbomer-, cellulose- and mineral oil-based artificial tear formulations. *Eur J Ophthalmol.* 2007;17(2):151-9. 26. McCann LC, Tomlinson A, Pearce EI, Papa V. Effectiveness of artificial tears in the management of evaporative dry eye. *Cornea.* 2012;31(1):1-5. 27. Rajendraprasad RM, Kwatra G, Batra N. Carboxymethyl cellulose versus hydroxypropyl methylcellulose tear substitutes for dry eye due to computer vision syndrome: comparison of efficacy and safety. *Int J Appl Basic Med Res.* 2021;11(1):4-8. 28. World report on vision. Geneva: World Health Organization; 2019 (<https://apps.who.int/iris/handle/10665/328717>, accessed 6 October 2023). 29. Safarzadeh M, Azizzadeh P, Akbarshahi P. Comparison of the clinical efficacy of preserved and preservative-free hydroxypropyl methylcellulose-dextran-containing eyedrops. *J Optom.* 2017;10(4):258-64. 30. Waduthantri S, Yong SS, Tan CH, Shen L, Lee MX, Nagarajan S, et al. Cost of dry eye treatment in an Asian clinic setting. *PLoS One.* 2012;7(6):e37711.