

		EMLc	Codes ATC: C03CA01
Indication	Heart failure	Code ICD11: BD0Z	
INN	Furosemide		
Type de médicament	Chemical agent		
Type de liste	Liste de base (EML) (EMLc)		
Additional notes	The square box does not apply to the listing of furosemide on the EMLc		
Formulations	Parenteral > General injections > IV: 10 mg per mL in 2 mL ampoule ; 10 mg per mL in 5 mL ampoule Oral > Liquid: 20 mg per 5 mL (EMLc) ; 50 mg per 5 mL (EMLc) Oral > Solid > tablet: 40 mg ; 20 mg		
Historique des statuts LME	Ajouté pour la première fois en 2005 (TRS 933) Modifié en 2007 (TRS 950) Modifié en 2021 (TRS 1035) Modifié en 2023 (TRS 1049)		
Sexe	Tous		
Âge	Aussi recommandé pour les enfants		
Équivalence thérapeutique	bumetanide (Codes ATC: C03CA02) torasemide (Codes ATC: C03CA04)		
Renseignements sur le brevet	Patents have expired in most jurisdictions Lire la suite sur les brevets . 		
Wikipédia	Furosemide 		
DrugBank	Furosemide 		

Résumé des preuves et recommandation du comité d'experts

Following the review of the age-appropriateness of formulations on the EMLc, the Expert Committee recommended the addition of new strength formulations of furosemide (injection 10 mg/mL in 5 mL; tablet 20 mg (EML and EMLc); oral liquid 50 mg/5 mL (EMLc). The Committee also recommended the removal of furosemide 10 mg tablets from the EMLc.

