

EMLc

Codes ATC: C03CA01

Indication	Oedema <span style="background-color: #008080; color: white; padding: 2px;">Code ICD11: MG29.Z</span>
INN	Furosemide
Type de médicament	Chemical agent
Type de liste	Liste de base (EML) (EMLc)
Additional notes	The square box applies only to the listing of furosemide on the EML.
Formulations	Parenteral > General injections > IV: 10 mg per mL in 2 mL ampoule ; 10 mg per mL in 5 mL ampoule Oral > Liquid: 20 mg per 5 mL (EMLc) ; 50 mg per 5 mL (EMLc) Oral > Solid > tablet: 40 mg ; 20 mg
Historique des statuts LME	Ajouté pour la première fois en 1977 (TRS 615) Modifié en 1979 (TRS 641) Modifié en 2007 (TRS 950) Modifié en 2021 (TRS 1035) Modifié en 2023 (TRS 1049)
Sexe	Tous
Âge	Aussi recommandé pour les enfants
Équivalence thérapeutique	bumetanide (Codes ATC: C03CA02) torasemide (Codes ATC: C03CA04)
Renseignements sur le brevet	Patents have expired in most jurisdictions Lire la suite <a href="#">sur les brevets.</a> 
Wikipédia	<a href="#">Furosemide</a> 
DrugBank	<a href="#">Furosemide</a> 

### Résumé des preuves et recommandation du comité d'experts

Following the review of the age-appropriateness of formulations on the EMLc, the Expert Committee recommended the addition of new strength formulations of furosemide (injection 10 mg/mL in 5 mL; tablet 20 mg (EML and EMLc); oral liquid 50 mg/5 mL (EMLc). The Committee also recommended the removal of furosemide 10 mg tablets from the EMLc.

