



		EMLc	Codes ATC: A06AA02
Indication	Palliative care	Code ICD11: QC7A	
INN	Docusate sodium		
Type de médicament	Chemical agent		
Type de liste	Liste de base (EML) (EMLc)		
Formulations	Oral > Liquid: 50 mg per 5 mL oral liquid Oral > Solid: 100 mg capsule		
Historique des statuts LME	Ajouté pour la première fois en 2009 (TRS 958) Modifié en 2013 (TRS 985)		
Sexe	Tous		
Âge	Aussi recommandé pour les enfants		
Équivalence thérapeutique	La recommandation concerne ce médicament spécifique		
Renseignements sur le brevet	Lire la suite sur les brevets. 		
Wikipédia	Docusate sodium 		
DrugBank	Docusate dodium (Docusate) 		

Résumé des preuves et recommandation du comité d'experts

Following a review of medicines for palliative care by the Expert Committee in 2013, docusate sodium was added to the EML for management of constipation in patients receiving palliative care. In 2013, the Expert Committee considered a review of medicines for palliative care in adults. To develop the list of proposed medicines, a working group was constituted by IAHP. Using WHO global mortality data this group identified the most common causes of death, and the most common and distressing symptoms in palliative care by means of a literature search. The final step was identification of the medicines to be recommended for the treatment of symptoms. Table 1 (page 49-50, TRS 985) lists the most common and distressing symptoms occurring in people receiving palliative care, and the medications included in the application. The Expert Committee noted that good-quality evidence for rational pharmacotherapy for many of the symptoms was lacking. The systematic reviews in the literature of symptom management related to patients with cancer. These reviews concluded that there was insufficient evidence to draw any firm conclusions on medicines for the treatment of symptoms such as fatigue, anxiety or anorexia in cancer. However, there is extensive experience in the use of these medicines for the treatment of most of the common symptoms experienced in terminally ill patients. The medicines that have been recommended in the application are already listed in the EML, under either palliative care or another indication. The unique position of the terminally ill patient and the objectives of providing palliative care means that experience needs to be taken into account. Given the above, much of the evidence for the efficacy of these drugs was based on studies that were not of patients receiving palliative care. The Expert Committee also noted the medicines that were included for palliative care in children. The Expert Committee recommended the listing of medicines for adults for common symptoms in palliative care, other than for pain, as shown in Table 2 (page 51-52, TRS 985). This list included docusate sodium for constipation. In 2013, medicines for palliative care were moved from Section 8.4 to Section 2.3 of the EML and EMLc - Medicines for other common symptoms in palliative care.

