




EMLc
ATC codes: N07AA02

Indication	Myasthenia gravis ICD11 code: 8C70.Z
INN	Pyridostigmine bromide
Medicine type	Chemical agent
List type	Complementary
Formulations	Parenteral > General injections > IV: 1 mg in 1 mL ampoule (pyridostigmine bromide) Oral > Solid: 60 mg (pyridostigmine bromide)
EML status history	First added in 1977 (TRS 615) Changed in 1979 (TRS 641) Changed in 1993 (TRS 850) Changed in 2003 (TRS 920) Changed in 2007 (TRS 950)
Sex	All
Age	Also recommended for children
Therapeutic alternatives	The recommendation is for this specific medicine
Patent information	Patents have expired in most jurisdictions Read more about patents . 
Wikipedia	Pyridostigmine 
DrugBank	Pyridostigmine 

Summary of evidence and Expert Committee recommendations

The EMLc Subcommittee noted that neostigmine metilsulfate injection was the preferred agent for the reversal of non-depolarising muscle block and endorsed neostigmine injection for inclusion in the EMLc. While neostigmine bromide tablets are licensed for use in infants and children for the treatment of myasthenia gravis, the Subcommittee noted that pyridostigmine has fewer cholinergic adverse effects and is now regarded as the first-line drug treatment for myasthenia gravis. The Subcommittee endorsed neostigmine (injection and tablet forms) and pyridostigmine (injection and tablet forms) for inclusion in the EMLc, with the latter retained in the Complementary List.

