# Pyridostigmine

Section: 20. Muscle relaxants (peripherally-acting) and cholinesterase inhibitors

EMLc | Codes ATC: N07AA02

| | |
|---|---|
| **Indication** | Myasthenia gravis Code ICD11: 8C60.Z |
| **INN** | Pyridostigmine bromide |
| **Type de médicament** | Chemical agent |
| **Type de liste** | Liste complémentaire |
| **Formulations** | **Parenteral > General injections > IV:** 5 mg per mL in ampoule or vial (pyridostigmine bromide)<br>**Oral > Solid > tablet:** 60 mg scored (pyridostigmine bromide) |
| **Historique des statuts LME** | Ajouté pour la première fois en 1977 (TRS 615)<br>Modifié en 1979 (TRS 641)<br>Modifié en 1993 (TRS 850)<br>Modifié en 2003 (TRS 920)<br>Modifié en 2007 (TRS 950)<br>Modifié en 2025 (TRS 1064) |
| **Sexe** | Tous |
| **Âge** | Aussi recommandé pour les enfants |
| **Équivalence thérapeutique** | La recommandation concerne ce médicament spécifique |
| **Renseignements sur le brevet** | Patents have expired in most jurisdictions<br>Lire la suite sur les brevets. |
| **Wikipédia** | Pyridostigmine |
| **DrugBank** | Pyridostigmine |

## Résumé des preuves et recommandation du comité d'experts

Following the review of the age-appropriateness of formulations on the EMLc, the Expert Committee recommended: - the deletion of pyridostigmine injection 1 mg in 1 mL ampoule; - the addition of pyridostigmine injection 5 mg/mL in ampoule or vial - to modify the formulation descriptions to better describe the available dosage forms.