Neostigmine

Section: 20. Muscle relaxants (peripherally-acting) and cholinesterase inhibitors

EMLc | Codes ATC: N07AA01

Indication	Anticholinesterase reversible Code ICD11: XM9D92
INN	Neostigmine bromide
Type de médicament	Chemical agent
Type de liste	Liste de base
Formulations	**Parenteral > General injections > unspecified:** 0.5 mg per mL in 1 mL ampoule (neostigmine metilsulfate) ; 2.5 mg per mL in 1 mL ampoule (neostigmine metilsulfate) **Oral > Solid > tablet:** 15 mg (neostigmine bromide) (EML)
Historique des statuts LME	Ajouté pour la première fois en 1977 (TRS 615) Modifié en 1979 (TRS 641) Modifié en 1989 (TRS 796) Modifié en 2003 (TRS 920) Modifié en 2007 (TRS 950) Modifié en 2025 (TRS 1064)
Sexe	Tous
Âge	Aussi recommandé pour les enfants
Équivalence thérapeutique	La recommandation concerne ce médicament spécifique
Renseignements sur le brevet	Patents have expired in most jurisdictions Lire la suite sur les brevets.
Wikipédia	Neostigmine
DrugBank	Neostigmine

Résumé des preuves et recommandation du comité d'experts

Following the review of the age-appropriateness of formulations on the EMLc, the Expert Committee recommended: - the deletion of neostigmine 15 mg tablets from the EMLc