




		EMLc	Codes ATC: S02AA10
Indication	Infectious diseases of external ear	Code ICD11: AA0Z	
Type de médicament	Chemical agent		
Type de liste	Liste de base (EML) (EMLc)		
Formulations	Local > Otological > drops: 2% in alcohol (EMLc)		
Historique des statuts LME	Ajouté pour la première fois en 2009 (TRS 958)		
Sexe	Tous		
Âge	Enfants (1 mois - 12 ans)		
Équivalence thérapeutique	La recommandation concerne ce médicament spécifique		
Renseignements sur le brevet	Patents have expired in most jurisdictions Lire la suite sur les brevets . 		
Wikipédia	Acetic acid 		
DrugBank	Acetic acid 		

Résumé des preuves et recommandation du comité d'experts

In 2009, the EMLc Subcommittee considered the review that was commissioned to identify essential medicines for the treatment of ear, nose and throat conditions in children. Based primarily on South African guidelines and WHO guidelines for treating ear, nose and throat conditions in children, the priority conditions identified were acute croup, epiglottitis, epistaxis, otitis externa, otitis media (acute and chronic), rhinosinusitis and sore throat. The Subcommittee noted that many medicines required for treating these conditions were already listed. However, several more needed to be considered for addition. These included preparations for both topical and systemic use. There was a lack of documented evidence for the efficacy and safety of most of the medicines that needed to be considered for addition. Most available data came from studies in adults or from those involving both adults and children. However, these medicines are recommended in widely accepted guidelines. The Subcommittee noted that there is evidence to show that antibiotic ear drops are of benefit in the treatment of otitis externa. There is evidence to suggest that quinolone ear drops are superior to other otic antimicrobial formulations. The Subcommittee considered the importance of the available information regarding combination antimicrobial agent–corticosteroid formulations for the treatment of otitis externa (1) and found no compelling evidence to support the inclusion of combination products. The Subcommittee therefore recommended that acetic acid ear drops and ciprofloxacin ear drops be added to the EMLc, the ciprofloxacin ear drops with a square box annotation. The Subcommittee decided that based on available evidence, the inclusion of a nasal corticosteroid could be recommended. Further, the Subcommittee recommended the inclusion of a decongestant nasal spray/nose drops, listing xylometazoline with a square box annotation. Topical ephedrine was not included in recognition of its potential for abuse. 1. Rosenfeld RM et al. Systematic review of topical antimicrobial therapy for acute otitis externa. *Otolaryngology and Head and Neck Surgery*, 2006, 134(4 Suppl):S24–48.

