



EMLc

Codes ATC: N02BA01

Indication	Mucocutaneous lymph node syndrome Code ICD11: 4A44.5
Type de médicament	Chemical agent
Type de liste	Liste complémentaire (EML) (EMLc)
Additional notes	Acute or chronic use for rheumatic fever, juvenile arthritis , Kawasaki disease
Formulations	Local > Rectal > Suppository: 150 mg ; 300 mg Oral > Solid > dispersible tablet: 75 mg ; 300 mg ; 500 mg Oral > Solid > tablet: 75 to 500 mg
Historique des statuts LME	Ajouté pour la première fois en 2007 (TRS 950) Modifié en 2013 (TRS 985) Modifié en 2025 (TRS 1064)
Sexe	Tous
Âge	Aussi recommandé pour les enfants
Équivalence thérapeutique	La recommandation concerne ce médicament spécifique
Renseignements sur le brevet	Patents have expired in most jurisdictions Lire la suite sur les brevets.
Wikipédia	Acetylsalicylic acid
DrugBank	Acetylsalicylic acid

Résumé des preuves et recommandation du comité d'experts

Following the review of the age-appropriateness of formulations of essential medicines for children, the Expert Committee recommended the following changes to the listing of acetylsalicylic acid: - the lower strength of the range of strengths for acetylsalicylic acid tablets be changed from 100 mg to 75 mg; - removal of the 50 mg strength of suppositories, and addition of 300 mg strength of suppositories; - addition of dispersible tablets (75 mg, 300 mg and 500 mg) to facilitate dosing in children unable to swallow solid tablets.

