




		<b>EMLc</b>	Codes ATC: <b>L01DB01</b>
<b>Indication</b>	Ewing sarcoma of bone and articular cartilage of unspecified sites	Code ICD11: <b>2C22.Z</b>	
<b>INN</b>	Doxorubicin		
<b>Type de médicament</b>	Chemical agent		
<b>Type de liste</b>	Liste complémentaire (EML) (EMLc)		
<b>Formulations</b>	Parenteral > General injections > IV: 10 mg in vial (hydrochloride) powder for injection ; 50 mg in vial (hydrochloride) powder for injection ; 2 mg per mL in 5 mL vial (hydrochloride) ; 2 mg per mL in 25 mL vial (hydrochloride)		
<b>Historique des statuts LME</b>	Ajouté pour la première fois en 2015 (TRS 994) Modifié en 2023 (TRS 1049)		
<b>Sexe</b>	Tous		
<b>Âge</b>	Aussi recommandé pour les enfants		
<b>Équivalence thérapeutique</b>	La recommandation concerne ce médicament spécifique		
<b>Renseignements sur le brevet</b>	Patents have expired in most jurisdictions Lire la suite <a href="#">sur les brevets.</a> 		
<b>Balises</b>	<span>Cancer</span>		
<b>Wikipédia</b>	<a href="#">Doxorubicin</a> 		
<b>DrugBank</b>	<a href="#">Doxorubicin</a> 		

## Résumé des preuves et recommandation du comité d'experts

Following the review of the age-appropriateness of formulations on the EMLc, the Expert Committee recommended the inclusion of an additional dose forms of doxorubicin (2 mg/mL (hydrochloride) in 5 mL, 25 mL vial) to the EML and EMLc.

[Rapport du comité d'experts](#) 

